




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/54, 9/10, 15/81, 15/82, 1/16, 5/10, A01N 27/067, C12P 7/64		A2	(11) International Publication Number: WO 00/60095
			(43) International Publication Date: 12 October 2000 (12.10.00)
(21) International Application Number: PCT/EP00/02701		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 28 March 2000 (28.03.00)		Published Without international search report and to be republished upon receipt of that report.	
(30) Priority Data: 99106656.4 1 April 1999 (01.04.99) EP 99111321.8 10 June 1999 (10.06.99) EP 60/180,687 7 February 2000 (07.02.00) US			
(71) Applicant (for all designated States except US): BASF PLANT SCIENCE GMBH [DE/DE]; D-67056 Ludwigshafen (DE).			
(72) Inventors; and (75) Inventors/Applicants (for US only): DAHLQVIST, Anders [SE/SE]; Hemmansvägen 2, S-244 66 Furulund (SE). STAHL, Ulf [SE/SE]; Liljegatan 7b, S-753 24 Uppsala (SE). LENMAN, Marit [SE/SE]; Revingsgatan 13a, S-223 59 Lund (SE). BANAS, Antoni [SE/PL]; Wiolinowa 14, PL-08110 Siedlce (PL). RONNE, Hans [SE/SE]; Dirigentvägen 169, S-756 54 Uppsala (SE). STYMNE, Sten [SE/SE]; Torrlösa 1380, S-269 90 Svalöv (SE).			
(74) Agent: FITZNER, Uwe; Lintorfer Str. 10, D-40878 Ratingen (DE).			

(54) Title: A NEW CLASS OF ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA MOLECULES ENCODING THESE ENZYMES



(57) Abstract

The present invention relates to the isolation, identification and characterization of nucleotide sequences encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol, to the said enzymes and a process for the production of triacylglycerols.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

A NEW CLASS OF ENZYMES IN THE BIOSYNTHETIC PATHWAY FOR THE
PRODUCTION OF TRIACYLGLYCEROL AND RECOMBINANT DNA
MOLECULES ENCODING THESE ENZYMES

- 5 The present invention relates to the isolation, identification and characterization of recombinant DNA molecules encoding enzymes catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.
- 10 Triacylglycerol (TAG) is the most common lipid-based energy reserve in nature. The main pathway for synthesis of TAG is believed to involve three sequential acyl-transfers from acyl-CoA to a glycerol backbone (1, 2). For many years, acyl-CoA : diacylglycerol acyltransferase (DAGAT), which catalyzes the third acyl transfer reaction, was thought to be the only unique enzyme involved in
- 15 TAG synthesis. It acts by diverting diacylglycerol (DAG) from membrane lipid synthesis into TAG (2). Genes encoding this enzyme were recently identified both in the mouse (3) and in plants (4, 5), and the encoded proteins were shown to be homologous to acyl-CoA : cholesterol acyltransferase (ACAT). It was also recently reported that another DAGAT exists in the oleaginous fungus
- 20 *Mortierella ramanniana*, which is unrelated to the mouse DAGAT, the ACAT gene family or to any other known gene (6).

The instant invention relates to novel type of enzymes and their encoding genes for transformation. More specifically, the invention relates to use of a

25 type of genes encoding a not previously described type of enzymes hereinafter designated phospholipid:diacylglycerol acyltransferases (PDAT), whereby this enzyme catalyses an acyl-CoA-independent reaction. The said type of genes expressed alone in transgenic organisms will enhance the total amount of oil (triacylglycerols) produced in the cells. The PDAT genes, in combination with a

30 gene for the synthesis of an uncommon fatty acid will, when expressed in transgenic organisms, enhance the levels of the uncommon fatty acids in the triacylglycerols.

There is considerable interest world-wide in producing chemical feedstock, such as fatty acids, for industrial use from renewable plant resources rather than non-renewable petrochemicals. This concept has broad appeal to
5 manufacturers and consumers on the basis of resource conservation and provides significant opportunity to develop new industrial crops for agriculture.

There is a diverse array of unusual fatty acids in oils from wild plant species and these have been well characterised. Many of these acids have industrial
10 potential and this has led to interest in domesticating relevant plant species to enable agricultural production of particular fatty acids.

Development in genetic engineering technologies combined with greater understanding of the biosynthesis of unusual fatty acids now makes it possible
15 to transfer genes coding for key enzymes involved in the synthesis of a particular fatty acid from a wild species into domesticated oilseed crops. In this way individual fatty acids can be produced in high purity and quantities at moderate costs.

20 In all crops like rape, sunflower, oilpalm etc., the oil (i.e. triacylglycerols) is the most valuable product of the seeds or fruits and other compounds like starch, protein, and fibre is regarded as by-products with less value. Enhancing the quantity of oil per weight basis at the expense of other compounds in oil crops would therefore increase the value of crop. If genes, regulating the allocation of
25 reduced carbon into the production of oil can be up-regulated, the cells will accumulate more oil on the expense of other products. Such genes might not only be used in already high oil producing cells, such as oil crops, but could also induce significant oil production in moderate or low oil containing crops such as e.g. soy, oat, maize, potato, sugarbeets, and turnips as well as in
30 micro-organisms.

Summary of the invention

Many of the unusual fatty acids of interest, e.g. medium chain fatty acids, hydroxy fatty acids, epoxy fatty acids and acetylenic fatty acids, have physical properties that are distinctly different from the common plant fatty acids. The present inventors have found that, in plant species naturally accumulating these uncommon fatty acids in their seed oil (i.e. triacylglycerol), these acids are absent, or present in very low amounts in the membrane (phospho)lipids of the seed. The low concentration of these acids in the membrane lipids is most likely a prerequisite for proper membrane function and thereby for proper cell functions. One aspect of the invention is that seeds of transgenic crops can be made to accumulate high amounts of uncommon fatty acids if these fatty acids are efficiently removed from the membrane lipids and channelled into seed triacylglycerols.

The inventors have identified a novel class of enzymes in plants catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the production of triacylglycerol through an acyl-CoA-independent reaction and that these enzymes (phospholipid:diacylglycerol acyltransferases, abbreviated as PDAT) are involved in the removal of hydroxylated, epoxygenated fatty acids, and probably also other uncommon fatty acids such as medium chain fatty acids, from phospholipids in plants.

This enzyme reaction was shown to be present in microsomal preparations from baker's yeast (*Saccharomyces cerevisiae*). The instant invention further pertains to an enzyme comprising an amino acid sequence as set forth in SEQ ID No. 2 or a functional fragment, derivate, allele, homolog or isoenzyme thereof. A so called 'knock out' yeast mutant, disrupted in the respective gene was obtained and microsomal membranes from the mutant was shown to totally lack PDAT activity. Thus, it was proved that the disrupted gene encodes a PDAT enzyme (SEQ ID NO. 1 and 2). Furtherm, this PDAT enzyme is

characterized through the amino acid sequence as set forth in SEQ ID NO 2 containing a lipase motif of the conserved sequence string FXKWVEA.

The instant invention pertains further to an enzyme comprising an amino acid sequence as set forth in SEQ ID NO. 1a, 2b or 5a or a functional fragment, 5 derivate, allele, homolog or isoenzyme thereof.

Further genes and/or proteins of so far unknown function were identified and are contemplated within the scope of the instant invention. A gene from *Schizosaccharomyces pombe*, SPBC776.14 (SEQ ID. NO. 3), a putative open reading frame CAA22887 of the SPBC776.14 (SEQ ID NO. 13) were identified.

10 Further *Arabidopsis thaliana* genomic sequences (SEQ ID NO. 4, 10 and 11) coding for putative proteins were identified, as well as a putative open reading frame AAC80628 from the *A. thaliana* locus AC 004557 (SEQ ID NO. 14) and a putative open reading frame AAD10668 from the *A. thaliana* locus AC 003027 (SEQ ID NO. 15) were identified.

15 Also, a partially sequenced cDNA clone from *Neurospora crassa* (SEQ ID NO. 9) and a *Zea mays* EST (Extended Sequence Tac) clone (SEQ ID NO. 7) and corresponding putative amino acid sequence (SEQ ID NO. 8) were identified. Finally, two cDNA clones were identified, one *Arabidopsis thaliana* EST (SEQ ID NO. 5 and corresponding predicted amino acid sequence SEQ ID NO. 6) 20 and a *Lycopersicon esculentum* EST clone (SEQ ID NO. 12) were identified.

Further, enzymes designated as PDAT comprising an amino acid sequence selected from the group consisting of sequences as set forth in SEQ ID NO 2a, 3a, 5b, 6 or 7b containing a lipase motif FXKWVEA are contemplated within the scope of the invention. Moreover, an enzyme comprising an amino acid 25 sequence encoded through a nucleotide sequence, a portion, derivate, allele or homolog thereof selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 or a functional fragment, derivate, allele, homolog or isoenzyme of the enzyme encoding amino acid sequence are included within the scope of the 30 invention.

A functional fragment of the instant enzyme is understood to be any polypeptide sequence which shows specific enzyme activity of a phospholipid:diacylglycerol acyltransferase (PDAT). The length of the functional fragment can for example vary in a range from about 660 ± 10 amino acids to 660 ± 250 amino acids, preferably from about 660 ± 50 to 660 ± 100 amino acids, whereby the „basic number“ of 660 amino acids corresponds in this case to the polypeptide chain of the PDAT enzyme of SEQ ID NO. 2 encoded by a nucleotide sequence according to SEQ ID NO. 1. Consequently, the „basic number“ of functional fulllength enzyme can vary in correspondance to the encoding nucleotide sequence.

A portion of the instant nucleotide sequence is meant to be any nucleotide sequence encoding a polypeptid which shows specific activity of a phospholipid:diacylglycerol acyltransferase (PDAT). The length of the nucleotide portion can vary in a wide range of about several hundreds of nucleotides based upon the coding region of the gene or a highly conserved sequence. For example the length varies in a range form about 1900 ± 10 to 1900 ± 1000 nucleotides, preferably form about 1900 ± 50 to 1900 ± 700 and more preferably form about 1900 ± 100 to 1900 ± 500 nucleotides. whereby the „basic number“ of 1900 nucleotides corresponds in this case to the encoding nucleotide sequence of the PDAT enzyme of SEQ ID NO. 1. Consequently, the „basic number“ of functional fulllength gene can vary.

An allelic variant of the instant nucleotide sequence is understood to be any different nucleotide sequence which encodes a polypeptide with a functionally equivalent function. The alleles pertain naturally occuring variants of the instant nucleotide sequences as well as synthetic nucleotide sequences produced by methods known in the art. Contemplated are even altered nucleotide sequences which result in an enzyme with altered activity and/or regulation or which is resistant against specific inhibitors. The instant invention further includes natural or synthetic mutations of the originally isolated nucleotide

sequences. These mutations can be substitution, addition, deletion, inversion or insertion of one or more nucleotides.

5 A homologous nucleotide sequence is understood to be a complementary sequence and/or a sequence which specifically hybridizes with the instant nucleotide sequence. Hybridizing sequences include similar sequences selected from the group of DNA or RNA which specifically interact to the instant nucleotide sequences under at least moderate stringency conditions which are known in the art. A preferred, non-limiting example of stringent hybridization
10 conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. This further includes short nucleotide sequences of e.g. 10 to 30 nucleotides, preferably 12 to 15 nucleotides. Included are also primer or hybridization probes.

15

A homologous nucleotide sequence included within the scope of the instant invention is a sequence which is at least about 40%, preferably at least about 50 % or 60%, and more preferably at least about 70%, 80% or 90% and most preferably at least about 95%, 96%, 97%, 98% or 99% or more homologous to
20 a nucleotide sequence of SEQ ID NO. 1.

All of the aforementioned definitions are true for amino acid sequences and functional enzymes and can easily transferred by a person skilled in the art.

Isoenzymes are understood to be enzymes which have the same or a similar
25 substrate specificity and/or catalytic activity but a different primary structure.

In a first embodiment, this invention is directed to nucleic acid sequences that encode a PDAT. This includes sequences that encode biologically active PDATs as well as sequences that are to be used as probes, vectors for
30 transformation or cloning intermediates. The PDAT encoding sequence may

encode a complete or partial sequence depending upon the intended use. All or a portion of the genomic sequence, cDNA sequence, precursor PDAT or mature PDAT is intended.

- 5 Further included is a nucleotide sequence selected from the group consisting of sequences set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 9b, 10, 10b or 11 or a portion, derivate, allele or homolog thereof. The invention pertains a partial nucleotide sequence corresponding to a fulllength nucleotide sequence selected from the group consisting of sequences set forth in SEQ ID No. 5, 5b, 6b, 7, 8b,
10 9, 11b or 12 or a portion, derivate, allele or homolog thereof. Moreover, a nucleotide sequence comprising a nucleotide sequence which is at least 40% homologous to a nucleotide sequence selected form the group consisting of those sequences set forth in SEQ ID No. 1 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 is contemplated within the scope of the invention.

15

The instant invention pertains to a gene construct comprising a said nucleotide sequences of the instant invention which is operably linked to a heterologous nucleic acid.

- The term operably linked means a serial organisation e.g. of a promotor, coding
20 sequence, terminator and/or further regulatory elements whereby each element can fulfill its original function during expression of the nucleotide sequence.

- Further, a vector comprising of a said nucleotide sequence of the instant invention is contemplated in the instant invention. This includes also an
25 expression vector as well as a vector further comprising a selectable marker gene and/or nucleotide sequences for the replication in a host cell and/or the integration into the genome of the host cell.

- In a different aspect, this invention relates to a method for producing a PDAT in
30 a host cell or progeny thereof, including genetically engineered oil seeds, yeast and moulds or any other oil accumulating organism, via the expression of a

construct in the cell. Cells containing a PDAT as a result of the production of the PDAT encoding sequence are also contemplated within the scope of the invention.

- 5 Further, the invention pertains a transgenic cell or organism containing a said nucleotide sequence and/or a said gene construct and/or a said vector. The object of the instant invention is further a transgenic cell or organism which is an eucaryotic cell or organism. Preferably, the transgenic cell or organism is a yeast cell or a plant cell or a plant. The instant invention further pertains said
- 10 transgenic cell or organism having an altered biosynthetic pathway for the production of triacylglycerol. A transgenic cell or organism having an altered oil content is also contemplated within the scope of this invention.

- Further, the invention pertains a transgenic cell or organism wherein the activity
- 15 of PDAT is altered in said cell or organism. This altered activity of PDAT is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme. Moreover, a transgenic cell or organism is included in the instant invention, wherein the altered biosynthetic pathway for the production of triacylglycerol is characterized by the prevention of
- 20 accumulation of undesirable fatty acids in the membrane lipids.

In a different embodiment, this invention also relates to methods of using a DNA sequence encoding a PDAT for increasing the oil-content within a cell.

- 25 Another aspect of the invention relates to the accommodation of high amounts of uncommon fatty acids in the triacylglycerol produced within a cell, by introducing a DNA sequence producing a PDAT that specifically removes these fatty acids from the membrane lipids of the cell and channel them into triacylglycerol. Plant cells having such a modification are also contemplated
- 30 herein.

Further, the invention pertains a process for the production of triacylglycerol, comprising growing a said transgenic cell or organism under conditions whereby the said nucleotide sequence is expressed and whereby the said transgenic cells comprising a said enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol forming triacylglycerol.

Moreover, triacylglycerols produced by the aforementioned process are included in scope of the instant invention.

Object of the instant invention is further the use of an instant nucleotide sequence and/or a said enzyme for the production of triacylglycerol and/or triacylglycerols with uncommon fatty acids. The use of a said instant nucleotide sequence and/or a said enzyme of the instant invention for the transformation of any cell or organism in order to be expressed in this cell or organism and result in an altered, preferably increased oil content of this cell or organism is also contemplated within the scope of the instant invention.

A PDAT of this invention includes any sequence of amino acids, such as a protein, polypeptide or peptide fragment obtainable from a microorganism, animal or plant source that demonstrates the ability to catalyse the production of triacylglycerol from a phospholipid and diacylglycerol under enzyme reactive conditions. By „enzyme reactive conditions“ is meant that any necessary conditions are available in an environment (e.g., such factors as temperature, pH, lack of inhibiting substances) which will permit the enzyme to function.

Other PDATs are obtainable from the specific sequences provided herein. Furthermore, it will be apparent that one can obtain natural and synthetic PDATs, including modified amino acid sequences and starting materials for synthetic-protein modelling from the exemplified PDATs and from PDATs which are obtained through the use of such exemplified sequences. Modified amino acid sequences include sequences that have been mutated, truncated,

increased and the like, whether such sequences were partially or wholly synthesised. Sequences that are actually purified from plant preparations or are identical or encode identical proteins thereto, regardless of the method used to obtain the protein or sequence, are equally considered naturally
5 derived.

10

Further, the nucleic acid probes (DNA and RNA) of the present invention can be used to screen and recover „homologous“ or „related“ PDATs from a variety of plant and microbial sources.

15

Further, it is also apparent that a person skilled in the art can, with the information provided in this application, in any organism identify a PDAT activity, purify an enzyme with this activity and thereby identify a „non-homologous“ nucleic acid sequence encoding such an enzyme.

The present invention can be essentially characterized by the following aspects:

1. Use of a PDAT gene (genomic clone or cDNA) for transformation.
- 20 2. Use of a DNA molecule according to item 1 wherein said DNA is used for transformation of any organism in order to be expressed in this organism and result in an active recombinant PDAT enzyme in order to increase oil content of the organism.
3. Use of a DNA molecule of item 1 wherein said DNA is used for
25 transformation of any organism in order to prevent the accumulation of undesirable fatty acids in the membrane lipids.
4. Use according to item 1, wherein said PDAT gene is used for transforming transgenic oil accumulating organisms engineered to produce any
30 uncommon fatty acid which is harmful if present in high amounts in membrane lipids, such as medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.

5. Use according to item 1, wherein said PDAT gene is used for transforming organisms, and wherein said organisms are crossed with other oil accumulating organisms engineered to produce any uncommon fatty acid which is harmful if present in high amounts in membrane lipids, comprising medium chain fatty acids, hydroxylated fatty acids, epoxygenated fatty acids and acetylenic fatty acids.
6. Use according to item 1, wherein the enzyme encoded by said PDAT gene or cDNA is coding for a PDAT with distinct acyl specificity.
7. Use according to item 1 wherein said PDAT encoding gene or cDNA, is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 30% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
8. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 40% or more *identical* to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
9. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 60% or more *identical* to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
10. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from *Saccharomyces cerevisiae*, or contain nucleotide sequences coding for an amino acid sequence 80% or more identical to the amino acid sequence of PDAT as presented in SEQ. ID. NO. 2.
11. Use according to item 1 wherein said PDAT encoding gene or cDNA is derived from plants or contain nucleotide sequences coding for an amino acid sequence 40% or more identical to the amino acid sequence of PDAT from *Arabidopsis thaliana* or to the protein encoded by the fulllength counterpart of the partial Zea mays, Lycopericon esculentum, or Neurospora crassa cDNA clones.

12. Transgenic oil accumulating organisms comprising, in their genome, a PDAT gene transferred by recombinant DNA technology or somatic hybridization.
13. Transgenic oil accumulating organisms according to item 12 comprising, in
5 their genome, a PDAT gene having specificity for substrates with a particular uncommon fatty acid and the gene for said uncommon fatty acid.
14. Transgenic organisms according to item 12 or 13 which are selected from the group consisting of fungi, plants and animals.
15. Transgenic organisms according to item 12 or 13 which are selected from
10 the group of agricultural plants.
16. Transgenic organisms according to item 12 or 13 which are selected from the group of agricultural plants and where said PDAT gene is expressed under the control of a storage organ specific promotor.
17. Transgenic organisms according to item 12 or 13 which are selected from
15 the group of agricultural plants and where said PDAT gene is expressed under the control of a seed promotor.
18. Oils from organisms according to item 12 – 17.
19. A method for altering acyl specificity of a PDAT by alteration of the nucleotide sequence of a naturally occurring encoding gene and as a
20 consequence of this alternation creating a gene encoding for an enzyme with novel acyl specificity.
20. A protein encoded by a DNA molecule according to item 1 or a functional fragment thereof.
21. A protein of item 20 designated phospholipid:diacylglycerol acyltransferase.
22. A protein of item 21 which has a distinct acyl specificity.
25
23. A protein of item 13 having the amino acid sequence as set forth in SEQ, ID NO. 2, 13, 14 or 15 (and the proteins encoded by the fulllength or partial genes set forth in SEQ. ID. NO. 1, 3, 4, 5, 7, 9, 10, 11 or 12) or an amino acid sequence with at least 30 % homology to said amino acid sequence.
- 30 24. A protein of item 23 isolated from *Saccharomyces cerevisiae*.

General methods:

Yeast strains and plasmids. The wild type yeast strains used were either FY1679 (*MAT α his3- Δ 200 leu2- Δ 1 trp1- Δ 6 ura3-52*) or W303-1A (*MAT α ADE2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1*) (7). The YNR008w::KanMX2 disruption strain FVKT004-04C(AL), which is congenic to FY1679, was obtained from the Euroscarf collection (8). A 2751 bp fragment containing the YNR008w gene with 583 bp of 5' and 183 bp of 3' flanking DNA was amplified from W303-1A genomic DNA using *Taq* polymerase with 5'-TCTCCATCTTCTGCAAAACCT-3' and 5'-CCTGTCAAAAACCTTCTCCTC-3' as primers. The resulting PCR product was purified by agarose gel electrophoresis and cloned into the *EcoRV* site of pBluescript (pbluescript-pdat). For complementation experiments, the cloned fragment was released from pBluescript by *HindIII*-*SacI* digestion and then cloned between the *HindIII* and *SacI* sites of pFL39 (9), thus generating pUS1. For overexpression of the PDAT gene, a 2202 bp *EcoRI* fragment from the pBluscript plasmid which contains only 24 bp of 5' flanking DNA was cloned into the *BamHI* site of the *GAL1-TPK2* expression vector pJN92 (12), thus generating pUS4.

Microsomal preparations. Microsomes from developing seeds of sunflower (*Helianthus annuus*), *Ricinus communis* and *Crepis palaestina* were prepared using the procedure of Stobart and Stymne (11). To obtain yeast microsomes, 1g of yeast cells (fresh weight) was re-suspended in 8 ml of ice-cold buffer (20 mM Tris-Cl, pH 7.9, 10 mM MgCl₂, 1 mM EDTA, 5 % (v/v) glycerol, 1 mM DTT, 0.3 M ammonium sulfate) in a 12 ml glass tube. To this tube, 4 ml of glass beads (diameter 0.45-0.5 mm) were added, and the tube was then heavily shaken (3 x 60 s) in an MSK cell homogenizer (B. Braun Melsungen AG, Germany). The homogenized suspension was centrifuged at 20,000 x g for 15 min at 6°C and the resulting supernatant was again centrifuged at 100,000 x g for 2 h at 6°C. The 100,000 x g pellet was resuspended in 0.1 M potassium

phosphate (pH 7.2), and stored at -80°C. It is subsequently referred to as the crude yeast microsomal fraction.

Lipid substrates. Radio-labeled ricinoleic (12-hydroxy-9-octadecenoic) and vernolic (12,13-epoxy-9-octadecenoic) acids were synthesized enzymatically from [1-¹⁴C]oleic acid and [1-¹⁴C]linoleic acid, respectively, by incubation with microsomal preparations from seeds of *Ricinus communis* and *Crepis palaestina*, respectively (12). The synthesis of phosphatidylcholines (PC) or phosphatidylethanolamines (PE) with ¹⁴C-labeled acyl groups in the *sn*-2 position was performed using either enzymatic (13), or synthetic (14) acylation of [¹⁴C]oleic, [¹⁴C]ricinoleic, or [¹⁴C]vernolic acid. Dioleoyl-PC that was labeled in the *sn*-1 position was synthesized from *sn*-1-[¹⁴C]oleoyl-lyso-PC and unlabeled oleic acid as described in (14). *Sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-DAG was synthesized from PC by the action of phospholipase C type XI from *B. Cereus* (Sigma Chemical Co.) as described in (15). Monovernoloyl- and divernoloyl-DAG were synthesized from TAG extracted from seeds of *Euphorbia lagascae*, using the TAG-lipase (*Rizhopus arrizus*, Sigma Chemical Co.) as previously described (16). Monoricinoleoyl-TAG was synthesized according to the same method using TAG extracted from Castor bean.

20

Lipid analysis. Total lipid composition of yeast were determined from cells harvested from a 40 ml liquid culture, broken in a glass-bead shaker and extracted into chloroform as described by Bligh and Dyer (17), and then separated by thin layer chromatography in hexane/diethylether/acetic acid (80:20:1) using pre-coated silica gel 60 plates (Merck). The lipid areas were located by brief exposure to I₂ vapors and identified by means of appropriate standards. Polar lipids, sterol-esters and triacylglycerols, as well as the remaining minor lipid classes, referred to as other lipids, were excised from the plates. Fatty acid methylesters were prepared by heating the dry excised material at 85 °C for 60 min in 2% (v/v) sulfuric acid in dry methanol. The methyl esters were extracted with hexane and analyzed by GLC through a 50 m

25
30

x 0.32 mm CP-Wax58-CB fused-silica column (Chrompack), with methylheptadecanoic acid as an internal standard. The fatty acid content of each fraction was quantified and used to calculate the relative amount of each lipid class. In order to determine the total lipid content, 3 ml aliquots from yeast
5 cultures were harvested by centrifugation and the resulting pellets were washed with distilled water and lyophilized. The weight of the dried cells was determined and the fatty acid content was quantified by GLC-analyses after conversion to methylesters as described above. The lipid content was then calculated as nmol fatty acid (FA) per mg dry weight yeast.

10 Enzyme assays. Aliquots of crude microsomal fractions (corresponding to 10 nmol of microsomal PC) from developing plant seeds or yeast cells were lyophilized over night. ¹⁴C-Labeled substrate lipids dissolved in benzene were then added to the dried microsomes. The benzene was evaporated under a
15 stream of N₂, leaving the lipids in direct contact with the membranes, and 0.1 ml of 50 mM potassium phosphate (pH 7.2) was added. The suspension was thoroughly mixed and incubated at 30°C for the time period indicated, up to 90 min. Lipids were extracted from the reaction mixture using chloroform and separated by thin layer chromatography in hexane/diethylether/acetic acid
20 (35:70:1.5) using silica gel 60 plates (Merck). The radioactive lipids were visualized and quantified on the plates by electronic autoradiography (Instant Imager, Packard, US).

Yeast cultivation. Yeast cells were grown at 28°C on a rotatory shaker in
25 liquid YPD medium (1% yeast extract, 2% peptone, 2% glucose), synthetic medium (18) containing 2% (v/v) glycerol and 2% (v/v) ethanol, or minimal medium (19) containing 16 g/l of glycerol.

The instant invention is further characterized by the following examples which
30 are not limiting:

Acyl-CoA-independent synthesis of TAG by oil seed microsomes. A large number of unusual fatty acids can be found in oil seeds (20). Many of these fatty acids, such as ricinoleic (21) and vernolic acids (22), are synthesized using phosphatidylcholin (PC) with oleoyl or linoleoyl groups esterified to the *sn*-2 position, respectively, as the immediate precursor. However, even though PC can be a substrate for unusual fatty acid synthesis and is the major membrane lipids in seeds, unusual fatty acids are rarely found in the membranes. Instead, they are mainly incorporated into the TAG. A mechanism for efficient and selective transfer of these unusual acyl groups from PC into TAG must therefore exist in oil seeds that accumulate such unusual fatty acids. This transfer reaction was biochemically characterized in seeds from castor bean (*Ricinus communis*) and *Crepis palaestina*, plants which accumulate high levels of ricinoleic and vernolic acid, respectively, and sunflower (*Helianthus annuus*), a plant which has only common fatty acids in its seed oil. Crude microsomal fractions from developing seeds were incubated with PC having ¹⁴C-labeled oleoyl, ricinoleoyl or vernoloyl groups at the *sn*-2 position. After the incubation, lipids were extracted and analyzed by thin layer chromatography. We found that the amount of radioactivity that was incorporated into the neutral lipid fraction increased linearly over a period of 4 hours (data not shown). The distribution of [¹⁴C]acyl groups within the neutral lipid fraction was analyzed after 80 min (Fig. 1). Interestingly the amount and distribution of radioactivity between different neutral lipids were strongly dependent both on the plant species and on the type of [¹⁴C]acyl chain. Thus, sunflower microsomes incorporated most of the label into DAG, regardless of the type of [¹⁴C]acyl group. In contrast, *R. communis* microsomes preferentially incorporated [¹⁴C]ricinoleoyl and [¹⁴C]vernoloyl groups into TAG, while [¹⁴C]oleyl groups mostly were found in DAG. *C. palaestina* microsomes, finally, incorporated only [¹⁴C]vernoloyl groups into TAG, with [¹⁴C]ricinoleyl groups being found mostly as free fatty acids, and [¹⁴C]oleyl groups in DAG. This shows that the high *in vivo* levels of ricinoleic acid and vernolic acid in the TAG pool of *R. communis*

and *C. palaestina*, respectively, can be explained by an efficient and selective transfer of the corresponding acyl groups from PC to TAG in these organisms.

The in-vitro synthesis of triacylglycerols in microsomal preparations of developing castor bean is summarized in table 1.

PDAT: a novel enzyme that catalyzes acyl-CoA independent synthesis of TAG. It was investigated if DAG could serve both as an acyl donor as well as an acyl acceptor in the reactions catalyzed by the oil seed microsomes. Therefore, unlabeled divernoloyl-DAG was incubated with either *sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-DAG or *sn*-1-oleoyl-*sn*-2-[¹⁴C]ricinoleoyl-PC in the presence of *R. communis* microsomes. The synthesis of TAG molecules containing both [¹⁴C]ricinoleoyl and vernoloyl groups was 5 fold higher when [¹⁴C]ricinoleoyl-PC served as acyl donor as compared to [¹⁴C]ricinoleoyl-DAG (fig.1B). These data strongly suggests that PC is the immediate acyl donor and DAG the acyl acceptor in the acyl-CoA-independent formation of TAG by oil seed microsomes. Therefore, this reaction is catalyzed by a new enzyme which we call phospholipid : diacylglycerol acyltransferase (PDAT).

PDAT activity in yeast microsomes. Wild type yeast cells were cultivated under conditions where TAG synthesis is induced. Microsomal membranes were prepared from these cells and incubated with *sn*-2-[¹⁴C]-ricinoleoyl-PC and DAG and the ¹⁴C-labeled products formed were analyzed. The PC-derived [¹⁴C]ricinoleoyl groups within the neutral lipid fraction mainly were found in free fatty acids or TAG, and also that the amount of TAG synthesized was dependent on the amount of DAG that was added to the reaction (Fig.2). The *in vitro* synthesis of TAG containing both ricinoleoyl and vernoloyl groups, a TAG species not present *in vivo*, from exogenous added *sn*-2-[¹⁴C]ricinoleoyl-PC and unlabelled vernoloyl-DAG (Fig. 2, lane 3) clearly demonstrates the existence of an acyl-CoA-independent synthesis of TAG involving PC and DAG as

substrates in yeast microsomal membranes. Consequently, TAG synthesis in yeast can be catalyzed by an enzyme similar to the PDAT found in plants.

The PDAT encoding gene in yeast.

5 A gene in the yeast genome (YNR008w) is known, but nothing is known about the function of YNR008w, except that the gene is not essential for growth under normal circumstances. Microsomal membranes were prepared from the yeast strain FVKT004-04C(AL) (8) in which this gene with unknown function had been disrupted. PDAT activity in the microsomes were assayed using PC with
10 radiolabelled fatty acids at the sn-2 position. The activity was found to be completely absent in the disruption strain (Fig. 2 lane 4). Significantly, the activity could be partially restored by the presence of YNR008w on the single copy plasmid pUS1 (Fig. 2 lane 5). Moreover, acyl groups of phosphatidylethanolamine (PE) were efficiently incorporated into TAG by
15 microsomes from the wild type strain whereas no incorporation occurred from this substrate in the mutant strain (data not shown). This shows that YNR008w encodes a yeast PDAT which catalyzes the transfer of an acyl group from the sn-2 position of phospholipids to DAG, thus forming TAG. It should be noted that no cholesterol esters were formed from radioactive PC even in incubations
20 with added ergosterols, nor were the amount of radioactive free fatty acids formed from PC affected by disruption of the YNR008w gene (data not shown). This demonstrates that yeast PDAT do not have cholesterol ester synthesising or phospholipase activities.

25 Increased TAG content in yeast cells that overexpress PDAT. The effect of overexpressing the PDAT-encoding gene was studied by transforming a wild type yeast strain with the pUS4 plasmid in which the gene is expressed from the galactose-induced *GAL1:TPK2* promoter. Cells containing the empty expression vector were used as a control. The cells were grown in synthetic
30 glycerol-ethanol medium, and expression of the gene was induced after either 2 hours (early log phase) or 25 hours (stationary phase) by the addition of

galactose. The cells were then incubated for another 21 hours, after which they were harvested and assays were performed. We found that overexpression of PDAT had no significant effect on the growth rate as determined by the optical density. However, the total lipid content, measured as μmol fatty acids per mg yeast dry weight, was 47% (log phase) or 29% (stationary phase) higher in the PDAT overexpressing strain than in the control. Furthermore, the polar lipid and sterolester content was unaffected by overexpression of PDAT. Instead, the elevated lipid content in these cells is entirely due to an increased TAG content (Fig. 3A,B). Thus, the amount of TAG was increased by 2-fold in PDAT overexpressing early log phase cells and by 40% in stationary phase cells. It is interesting to note that a significant increase in the TAG content was achieved by overexpressing PDAT even under conditions (*i.e.* in stationary phase) where DAGAT is induced and thus contributes significantly to TAG synthesis. *In vitro* PDAT activity assayed in microsomes from the PDAT overexpressing strain was 7-fold higher than in the control strain, a finding which is consistent with the increased levels of TAG that we observed *in vivo* (Fig. 3C). These results clearly demonstrate the potential use of the PDAT gene in increasing the oil content in transgenic organisms.

Substrate specificity of yeast PDAT. The substrate specificity of yeast PDAT was analyzed using microsomes prepared from the PDAT overexpressing strain (see Fig. 4). The rate of TAG synthesis, under conditions given in figure 4 with di-oleoyl-PC as the acyl-donor, was 0.15 nmol per min and mg protein. With both oleoyl groups of PC labeled it was possible, under the given assay conditions, to detect the transfer of 11 pmol/min of [^{14}C]oleoyl chain into TAG and the formation of 15 pmol/min of lyso-PC. In microsomes from the PDAT-deficient strain, no TAG at all and only trace amounts of lyso-PC was detected, strongly suggesting that yeast PDAT catalyses the formation of equimolar amounts of TAG and lyso-PC when supplied with PC and DAG as substrates. The fact that somewhat more lyso-PC than TAG is formed can be

explained by the presence of a phospholipase in yeast microsomes, which produces lyso-PC and unesterified fatty acids from PC.

The specificity of yeast PDAT for different acyl group positions was investigated by incubating the microsomes with di-oleoyl-PC carrying a $[^{14}\text{C}]$ acyl group either at the *sn*-1 position (Fig. 4A bar 2) or the *sn*-2 position (Fig. 4A bar 3). We found that the major ^{14}C -labeled product formed in the former case was lyso-PC, and in the latter case TAG. We conclude that yeast PDAT has a specificity for the transfer of acyl groups from the *sn*-2 position of the phospholipid to DAG, thus forming *sn*-1-lyso-PC and TAG. Under the given assay conditions, trace amounts of ^{14}C -labelled DAG is formed from the *sn*-1 labeled PC by the reversible action of a CDP-choline : choline phosphotransferase. This labeled DAG can then be further converted into TAG by the PDAT activity. It is therefore not possible to distinguish whether the minor amounts of labeled TAG that is formed in the presence of di-oleoyl-PC carrying a $[^{14}\text{C}]$ acyl group in the *sn*-1 position, is synthesized directly from the *sn*-1-labeled PC by a PDAT that also can act on the *sn*-1 position, or if it is first converted to *sn*-1-labeled DAG and then acylated by a PDAT with strict selectivity for the transfer of acyl groups at the *sn*-2 position of PC. Taken together, this shows that the PDAT encoded by YNR008w catalyses an acyl transfer from the *sn*-2 position of PC to DAG, thus causing the formation of TAG and lyso-PC.

The substrate specificity of yeast PDAT was further analyzed with respect to the headgroup of the acyl donor, the acyl group transferred and the acyl chains of the acceptor DAG molecule. The two major membrane lipids of *S. cerevisiae* are PC and PE, and as shown in Fig. 4B (bars 1 and 2), dioleoyl-PE is nearly 4-fold more efficient than dioleoyl-PC as acyl donor in the PDAT-catalyzed reaction. Moreover, the rate of acyl transfer is strongly dependent on the type of acyl group that is transferred. Thus, a ricinoleoyl group at the *sn*-2 position of PC is 2.5 times more efficiently transferred into TAG than an oleoyl

group in the same position (Fig. 4B bars 1 and 3). In contrast, yeast PDAT has no preference for the transfer of vernoloyl groups over oleoyl groups (Fig. 4B bars 1 and 4). The acyl chain of the acceptor DAG molecule also affects the efficiency of the reaction. Thus, DAG with a ricinoleoyl or a vernoloyl group is a more efficient acyl acceptor than dioleoyl-DAG (Fig. 4B bars 1, 5 and 6). Taken together, these results clearly show that the efficiency of the PDAT-catalyzed acyl transfer is strongly dependent on the properties of the substrate lipids.

PDAT genes. Nucleotide and amino acid sequences of several PDAT genes are given as SEQ ID No. 1 through 15. Further provisional and/or partial sequences are given as SEQ ID NO 1a through 5a and 1b through 11b, respectively. One of the Arabidopsis genomic sequences (SEQ ID NO. 4) identified an Arabidopsis EST cDNA clone; T04806. This cDNA clone was fully characterised and the nucleotide sequence is given as SEQ ID NO. 5. Based on the sequence homology of the T04806 cDNA and the *Arabidopsis thaliana* genomic DNA sequence (SEQ ID NO 4) it is apparent that an additional A is present at position 417 in the cDNA clone (data not shown). Excluding this nucleotide would give the amino acid sequence depicted in SEQ ID NO. 12.

Increased TAG content in seeds of Arabidopsis thaliana that express the yeast PDAT. For the expression of the yeast PDAT gene in *Arabidopsis thaliana* an EcoRI fragment from the pBluescript-PDAT was cloned together with napin promotor (25) into the vector pGPTV-KAN (26). A plasmid (pGNapPDAT) having the yeast PDAT gene in the correct orientation was identified and transformed into *Agrobacterium tumefaciens*. These bacteria were used to transform *Arabidopsis thaliana* columbia (C-24) plants using the root transformation method (27). Plants transformed with an empty vector were used as controls.

First generation seeds (T1) were harvested and germinated on kanamycin containing medium. Second generation seeds (T2) were pooled from individual plants and their fatty acid contents analysed by quantification of their methyl

esters by gas liquid chromatography after methylation of the seeds with 2% sulphuric acid in methanol at 85 °C for 1,5 hours. Quantification was done with heptadecanoic acid methyl esters as internal standard.

5 From the transformation with pGNapPDAT one T1 plant (26-14) gave raise to seven T2 plants of which 3 plants yielded seeds with statistically (in a mean difference two-sided test) higher oil content than seeds from T2 plants generated from T1 plant 32-4 transformed with an empty vector (table 2).

References cited in the description:

1. Bell, R. M. & Coleman, R. A. (1980) *Annu. Rev. Biochem.* **49**, 459-487.
2. Stymne, S. & Stobart, K. (1987) in *The biochemistry of plants: a comprehensive treatise*, Vol. 9, ed. Stumpf, P. K. (Academic Press, New York), pp. 175-214.
3. Cases, S. et al. (1998) *Proc. Natl. Acad. Sci. U S A* **95**, 13018-13023.
4. Hobbs, D. H., Lu, C. & Hills, M. J. (1999) *FEBS Lett.* **452**, 145-9
5. Zou, J., Wei, Y., Jako, C., Kumar, A., Selvaraj, G. & Taylor, D. C. (1999) *Plant J.* **19**, 645-653.
6. Lardizabal, K., Hawkins, D., Mai, J., & Wagner, N. (1999) Abstract presented at the Biochem. Mol. Plant Fatty Acids Glycerolipids Symposium, South Lake Tahoe, USA.
7. Thomas, B. J. & Rothstein, R. (1989) *Cell* **56**, 619-630.
8. Entian, K.-D. & Kötter, P. (1998) *Meth. Microbiol.* **26**, 431-449.
9. Kern, L., de Montigny, J., Jund, R. & Lacroute, F. (1990) *Gene* **88**, 149-157.
10. Ronne, H., Carlberg, M., Hu, G.-Z. & Nehlin, J. O. (1991) *Mol. Cell. Biol.* **11**, 4876-4884.
11. Stobart, K. & Stymne, S. (1990) in *Method in Plant Biochemistry*, vol 4, eds. Harwood, J. L. & Bowyer, J. R. (Academic press, London), pp. 19-46.
12. Bafor, M., Smith, M. A., Jonsson, L., Stobrt, A. K. & Stymne, S. (1991) *Biochem. J.* **280**, 507-514.
13. Banas, A., Johansson, I. & Stymne, S. (1992) *Plant Science* **84**, 137-144.
14. Kanda, P. & Wells, M. A. (1981) *J. Lipid. Res.* **22**, 877-879.
15. Ståhl, U., Ek, B. & Stymne, S. (1998) *Plant Physiol.* **117**, 197-205.
16. Stobart, K., Mancha, M. & Lenman M, Dahlqvist, A. & Stymne, S. (1997) *Planta* **203**, 58-66.
17. Bligh, E. G. & Dyer, W. J. (1959) *Can. J. Biochem. Physiol.* **37**, 911-917.
18. Sherman, F., Fink, G. R. & Hicks, J. B. (1986) in *Laboratory Course Manual for Methods in Yeast Genetics* (Cold Spring Harbor Laboratory)
19. Meesters, P. A. E. P., Huijberts, G. N. M. and Eggink, G. (1996) *Appl. Microbiol. Biotechnol.* **45**, 575-579.
20. van de Loo, F. J., Fox, B. G. & Sommerville, C. (1993), in *Lipid metabolism in plants*, ed. Moore, T. S. (CRC Press, Inc.), pp. 91-126.
21. van de Loo, F. J., Broun, P., Turner, S. & Sommerville, S. (1995) *Proc. Natl.*

Acad. Sci. U S A **95**, 6743-6747.

22. Lee, M., Lenman, M., Banas, A., Bafor, M., Singh, S., Schweizer, M., Nilsson, R., Liljenberg, C., Dahlqvist, A., Gummesson, P-O., Sjö Dahl, S., Green, A., and Stymne, S. (1998) *Science* **280**, 915-918.

5 23. Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997) *Nucl. Acids Res.* **24**, 4876-4882.

24. Saitou, N. & Nei, M. (1987) *Mol. Biol. Evol.* **4**, 406-425.

25. Stålberg, K., Ellerström, M., Josefsson, L., & Rask, L. (1993) *Plant Mol. Biol.* **23**, 671

10 26. Becker, D., Kemper, E., Schell, J., Masterson, R. (1992) *Plant Mol. Biol.* **20**, 1195

27. D. Valvekens, M. Van Montagu, and Van Lusbettens (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 5536

15

Description of Figures

FIG. 1.

Metabolism of ^{14}C -labeled PC into the neutral lipid fraction by plant

5 microsomes. (A) Microsomes from developing seeds of sunflower, *R. communis* and *C. palaestina* were incubated for 80 min at 30°C with PC (8 nmol) having oleic acid in its *sn*-1 position, and either ^{14}C -labeled oleic, ricinoleic or vernolic acid in its *sn*-2 position. Radioactivity incorporated in TAG (open bars), DAG (solid bars), and unsterified fatty acids (hatched bars) was
10 quantified using thin layer chromatography followed by electronic autoradiography, and is shown as percentage of added labeled substrate. (B) Synthesis *in vitro* of TAG carrying two vernoloyl and one [^{14}C]ricinoleoyl group by microsomes from *R. communis*. The substrates added were unlabeled divernoloyl-DAG (5 nmol), together with either *sn*-1-oleoyl-*sn*-2-[^{14}C]ricinoleoyl-DAG (0.4 nmol, 7700 dpm/nmol) or *sn*-1-oleoyl-*sn*-2-[^{14}C]ricinoleoyl-PC (0.4
15 nmol, 7700 dpm/nmol). The microsomes were incubated with the substrates for 30 min at 30°C, after which samples were removed for lipid analysis as described in the section „general methods“. The data shown are the average of two experiments.

20

FIG. 2.

PDAT activity in yeast microsomes, as visualized by autoradiogram of neutral lipid products separated on TLC. Microsomal membranes (10 nmol of PC) from

the wild type yeast strain FY1679 (lanes 1-3), a congenic yeast strain
25 (FVKT004-04C(AL)) that is disrupted for YNR008w (lane 4) or the same disruption strain transformed with the plasmid pUS1, containing the YNR008w gene behind its native promotor (lane 5), were assayed for PDAT activity. As substrates, we used 2 nmol *sn*-1-oleoyl-*sn*-2-[^{14}C]ricinoleoyl-PC together with either 5 nmol of dioleoyl-DAG (lanes 2, 4 and 5) or *rac*-oleoyl-vernoleoyl-DAG
30 (lane 3). The enzymatic assay and lipid analysis was performed as described in Materials and Methods. The cells were precultured for 20 h in liquid YPD

medium, harvested and re-suspended in an equal volume of minimal medium (19) containing 16 g/l glycerol. The cells were then grown for an additional 24 h prior to being harvested. Selection for the plasmid was maintained by growing the transformed cells in synthetic medium lacking uracil (18). Abbreviations: 1-OH-TAG, monoricinoleoyl-TAG; 1-OH-1-ep-TAG, monoricinoleoyl-monovernoloyl-TAG; OH-FA, unesterified ricinoleic acid.

Fig. 3.

Lipid content (A,B) and PDAT activity (C) in PDAT overexpressing yeast cells.

- 10 The PDAT gene in the plasmid pUS4 was overexpressed from the galactose-induced *GAL1-TPK2* promotor in the wild type strain W303-1A (7). Its expression was induced after (A) 2 hours or (B) 25 hours of growth by the addition of 2% final concentration (w/v) of galactose. The cells were then incubated for another 22 hours before being harvested. The amount of lipids of
- 15 the harvested cells was determined by GLC-analysis of its fatty acid contents and is presented as μmol fatty acids per mg dry weight in either TAG (open bar), polar lipids (hatched bar), sterol esters (solid bar) and other lipids (striped bar). The data shown are the mean values of results with three independent yeast cultures. (C) *In vitro* synthesis of TAG by microsomes prepared from
- 20 yeast cells containing either the empty vector (vector) or the PDAT plasmid (+ PDAT). The cells were grown as in Fig. 3A. The substrate lipids dioleoyl-DAG (2.5 nmol) and *sn*-1-oleoyl-*sn*-2-[^{14}C]-oleoyl-PC (2 nmol) were added to aliquots of microsomes (10 nmol PC), which were then incubated for 10 min at 28 °C. The amount of label incorporated into TAG was quantified by electronic
- 25 autoradiography. The results shown are the mean values of two experiments.

FIG. 4.

- Substrate specificity of yeast PDAT. The PDAT activity was assayed by incubating aliquots of lyophilized microsomes (10 nmol PC) with substrate lipids
- 30 at 30°C for 10 min (panel A) or 90 min (panel B). Unlabeled DAG (2.5 nmol) was used as substrates together with different labeled phospholipids, as shown

in the figure. (A) *Sn*-position specificity of yeast PDAT regarding the acyl donor substrate. Dioleoyl-DAG together with either *sn*-1- ^{14}C -oleoyl-*sn*-2- ^{14}C -oleoyl-PC (di- ^{14}C -PC), *sn*-1- ^{14}C -oleoyl-*sn*-2-oleoyl-PC (*sn*1- ^{14}C -PC) or *sn*-1-oleoyl-*sn*-2- ^{14}C -oleoyl-PC (*sn*2- ^{14}C -PC). (B) Specificity of yeast PDAT regarding phospholipid headgroup and of the acyl composition of the phospholipid as well as of the diacylglycerol. Dioleoyl-DAG together with either *sn*-1-oleoyl-*sn*-2- ^{14}C -oleoyl-PC (oleoyl-PC), *sn*-1-oleoyl-*sn*-2- ^{14}C -oleoyl-PE (oleoyl-PE), *sn*-1-oleoyl-*sn*-2- ^{14}C -ricinoleoyl-PC (ricinoleoyl-PC) or *sn*-1-oleoyl-*sn*-2- ^{14}C -vernoloyl-PC (vernoloyl-PC). In the experiments presented in the 2 bars to the far right, monoricinoleoyl-DAG (ricinoleoyl-DAG or mono-vernoloyl-DAG (vernoloyl-DAG) were used together with *sn*-1-oleoyl-*sn*-2- ^{14}C -oleoyl-PC. The label that was incorporated into TAG (solid bars) and lyso-PC (LPC, open bars) was quantified by electronic autoradiography. The results shown are the mean values of two experiments. The microsomes used were from W303-1A cells overexpressing the PDAT gene from the *GAL1-TPK2* promotor, as described in Fig. 3. The expression was induced at early stationary phase and the cells were harvested after an additional 24 h.

TAB.1:

In vitro synthesis of triacylglycerols in microsomal preparations of developing castor bean. Aliquots of microsomes (20 nmol PC) were lyophilised and substrate lipids were added in benzene solution: (A) 0.4 nmol ^{14}C -DAG (7760 dpm/nmol) and where indicated 1.6 nmol unlabelled DAG; (B) 0.4 nmol ^{14}C -DAG (7760 dpm/nmol) and 5 nmol unlabelled di-ricinoleoyl-PC and (C) 0.25 nmol ^{14}C -PC (4000 dpm/nmol) and 5 nmol unlabelled DAG. The benzene was evaporated by N_2 and 0.1 ml of 50 mM potassium phosphate was added, thoroughly mixed and incubated at 30 °C for (A) 20 min.; (B) and (C) 30 min.. Assays were terminated by extraction of the lipids in chloroform. The lipids were then separated by thin layer chromatography on silica gel 60 plates

(Merck; Darmstadt, Germany) in hexan/diethylether/acetic 35:70:1.5. The radioactive lipids were visualised and the radioactivity quantified on the plate by electronic autoradiography (Instant Imager, Packard, US). Results are presented as mean values of two experiments.

5

Radioactivity in different triacylglycerols (TAG) species formed. Abbreviations used: 1-OH-, mono-ricinoleoyl-; 2-OH, di-ricinoleoyl-; 3-OH-, triricinoleoyl; 1-OH-1-ver-, mono-ricinoleoyl-monovernoleoyl-; 1-OH-2-ver-, mono-ricinoleoyl-divernoleoyl-. Radiolabelled DAG and PC were prepared enzymatically. The radiolabelled ricinoleoyl group is attached at the sn-2-position of the lipid and unlabelled oleoyl group at the sn-1-position. Unlabelled DAG with vernoleoyl- or ricinoleoyl chains were prepared by the action of TAG lipase (6) on oil of *Euphorbia lagascae* or Castor bean, respectively. Synthetic di-ricinoleoyl-PC was kindly provided from Metapontum Agribios (Italy).

15

TAB.2:

Total fatty acids per mg of T2 seeds pooled from individual *Arabidopsis thaliana* plants transformed with yeast PDAT gene under the control of napin promotor (26-14) or transformed with empty vector (32-4).

20

* = statistical difference between control plants and PDAT transformed plants in a mean difference two-sided test at $\alpha = 5$.

Description of the SEQ ID:

5 *SEQ ID NO. 1:* Genomic DNA sequence and suggested amino acid sequence of the *Saccharomyces cerevisiae* PDAT gene, YNR008w, with GenBank accession number Z71623 and Y13139, and with nucleotide ID number 1302481.

SEQ ID NO. 2: The amino acid sequence of the suggested open reading frame YNR008w from *Saccharomyces cerevisiae*.

10 *SEQ ID NO. 3:* Genomic DNA sequence of the *Schizosaccharomyces pombe* gene SPBC776.14.

SEQ ID NO. 4: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AB006704.

15 *SEQ ID NO. 5:* Nucleotide sequence of the *Arabidopsis thaliana* cDNA clone with GenBank accession number T04806, and nucleotide ID number 315966.

SEQ ID NO. 6: Predicted amino acid sequence of the *Arabidopsis thaliana* cDNA
20 clone with GenBank accession number T04806.

SEQ ID NO. 7: Nucleotide and amino acid sequence of the *Zea mays* EST clone with GenBank accession number AI491339, and nucleotide ID number 4388167.

25 *SEQ ID NO. 8:* Predicted amino acid sequence of the *Zea mays* EST clone with GenBank accession number AI491339, and nucleotide ID number 4388167.

SEQ ID NO. 9: DNA sequence of part of the *Neurospora crassa* EST clone W07G1, with GenBank accession number AI398644, and nucleotide ID number
30 4241729.

SEQ ID NO. 10: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AC004557.

SEQ ID NO. 11: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with GenBank accession number AC003027.

SEQ ID NO. 12: DNA sequence of part of the *Lycopersicon esculentum* cDNA clone with GenBank accession number AI486635.

SEQ ID NO. 13: Amino acid sequence of the *Schizosaccharomyces pombe* putative open reading frame CAA22887 of the *Schizosaccharomyces pombe* gene SPBC776.14.

SEQ ID NO. 14: Amino acid sequence of the *Arabidopsis thaliana* putative open reading frame AAC80628 derived from the *Arabidopsis thaliana* locus with GenBank accession number AC004557.

SEQ ID NO 15: Amino acid sequence of the *Arabidopsis thaliana* putative open reading frame AAD10668 derived from the *Arabidopsis thaliana* locus with GenBank accession number AC003027.

Further provisional and/or partial sequences are defined through the following SEQ IDs:

SEQ ID NO. 1a: The amino acid sequence of the yeast ORF YNR008w from *Saccharomyces cerevisiae*.

SEQ ID NO. 2a: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AC004557).

SEQ ID NO. 3a: Amino acid sequence of the region of the *Arabidopsis thaliana* genomic sequence (AB006704).

5 SEQ ID NO. 4a: The corresponding genomic DNA sequence and amino acid sequence of the yeast ORF YNROO8w from *Saccharomyces cerevisiae*.

10 SEQ ID NO. 5a: The amino acid sequence of the yeast ORF YNROO8w from *Saccharomyces cerevisiae* derived from the corresponding genomic DNA sequence.

15 SEQ ID NO. 1b: Genomic DNA sequence of the *Saccharomyces cerevisiae* PDAT gene, YNR008w, genebank nucleotide ID number 1302481, and the suggested YNR008w amino acid sequence.

SEQ ID NO. 2b: The suggested amino acid sequence of the yeast gene YNR008w from *Saccharomyces cerevisiae*.

20 SEQ ID NO. 3b: Genomic DNA sequence of the *Schizosaccharomyces pombe* gene SPBC776.14.

SEQ ID NO. 4b: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AB006704.

25 SEQ ID NO. 5b: Nucleotide sequence and the corresponding amino acid sequence of the *Arabidopsis thaliana* EST-clone with genebank accession number T04806, and ID number 315966.

30 SEQ ID NO. 6b: Nucleotide and amino acid sequence of the *Zea mays* cDNA clone with genebank ID number 4388167.

SEQ ID NO. 7b: Amino acid sequence of the *Zea mays* cDNA clone with genebank ID number 4388167.

5 *SEQ ID NO. 8b*: DNA sequence of part of the *Neurospora crassa* cDNA clone WO7G1, ID number 4241729.

SEQ ID NO. 9b: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AC004557.

10 *SEQ ID NO. 10b*: Genomic DNA sequence of part of the *Arabidopsis thaliana* locus with genebank accession number AC003027.

SEQ ID NO. 11b: DNA sequence of part of the *Lycopersicon esculentum* cDNA clone with genebank accession number AI486635.

15

20

25

30

Claims

1. An enzyme catalysing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway
5 for the production of triacylglycerol.
2. An enzyme according to claim 1, comprising an amino acid sequence as set forth in SEQ ID No. 2 or a functional fragment, derivate, allele, homolog or isoenzyme thereof.
10
3. An enzyme according to claims 1 or 2 designated as phospholipid:diacylglycerol acyltransferase (PDAT).
4. An enzyme according to claims 1 to 3, comprising an amino acid sequence
15 as set forth in SEQ ID No. 1a, 2b or 5a or a functional fragment, derivate, allele, homolog or isoenzyme thereof.
5. An enzyme according to claims 1 to 4, comprising an amino acid sequence selected from the group consisting of sequences as set forth in SEQ ID No.
20 2a, 3a, 5b, 6, 7b, 8, 13, 14 or 15 or a functional fragment, derivate, allele, homolog or isoenzyme thereof.
6. An enzyme according to claims 1 to 5, comprising an amino acid sequence encoded through a nucleotide sequence, a portion, derivate, allele or
25 homolog thereof selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12 or a functional fragment, derivate, allele, homolog or isoenzyme of the enzyme encoding amino acid sequence.
- 30 7. A nucleotide sequence encoding an enzyme catalysing in an acyl-CoA-independent reaction the transfer of fatty acids from phospholipids to

diacylglycerol in the biosynthetic pathway for the production of triacylglycerol.

- 5 8. A nucleotide sequence according to claim 7 encoding an enzyme designated as phospholipid:diacylglycerol acyltransferase (PDAT).
9. A nucleotide sequence according to claims 7 or 8, selected from the group consisting of sequences as set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 9b, 10, 10b or 11 or a portion, derivate, allele or homolog thereof.
- 10 10. A partial nucleotide sequence corresponding to a fulllength nucleotide sequence according to claims 7 to 9, selected from the group consisting of sequences as set forth in SEQ ID No. 5, 5b, 6b, 7, 8b, 9, 11b or 12 or a portion, derivate, allele or homolog thereof.
- 15 11. A nucleotide sequence according to claims 7 to 10, comprising a nucleotide sequence which is at least 40% homologous to a nucleotide sequence selected from the group consisting of those sequences set forth in SEQ ID No. 1, 1b, 3, 3b, 4, 4a, 4b, 5, 5b, 6b, 7, 8b, 9, 9b, 10, 10b, 11, 11b or 12.
- 20 12. A gene construct comprising a nucleotide sequence according to claims 7 to 11 operably linked to a heterologous nucleic acid.
13. A vector comprising a nucleotide sequence according to claims 7 to 11 or a gene construct according to claim 12.
- 25 14. A vector according to claim 13, which is an expression vector.
15. A vector according to claims 13 or 14, further comprising a selectable marker gene and/or nucleotide sequences for the replication in a host cell or the integration into the genome of the host cell.
- 30

16. A transgenic cell or organism containing a nucleotide sequence according to claims 7 to 11 and/or a gene construct according to claim 12 and/or a vector according to claims 13 to 15.

5

17. A transgenic cell or organism according to claim 16 which is an eucaryotic cell or organism.

18. A transgenic cell or organism according to claims 16 or 17 which is a yeast cell or a plant cell or a plant.

10

19. A transgenic cell or organism according to claims 16 to 18 having an altered biosynthetic pathway for the production of triacylglycerol.

20. A transgenic cell or organism according to claims 16 to 19 having an altered oil content.

15

21. A transgenic cell or organism according to claims 16 to 20 wherein the activity of PDAT is altered.

20

22. A transgenic cell or organism according to claims 16 to 21 wherein the altered activity of PDAT is characterized by an alteration in gene expression, catalytic activity and/or regulation of activity of the enzyme.

23. A transgenic cell or organism according to claims 16 to 22 wherein the altered biosynthetic pathway for the production of triacylglycerol is characterized by the prevention of accumulation of undesirable fatty acids in the membrane lipids.

25

24. A process for the production of triacylglycerol, comprising growing a transgenic cell or organism according to claims 16 to 23 under conditions

30

whereby the said nucleotide sequence according to claims 7 to 11 is expressed.

25. Triacylglycerols produced by a process according to claim 24.

5

26. Use of a nucleotide sequence according to claims 7 to 11 and/or an enzyme according to claims 1 to 6 for the production of triacylglycerol and/or triacylglycerols with uncommon fatty acids.

10 27. Use of a nucleotide sequence according to claims 7 to 11 and/or an enzyme according to claims 1 to 6 for the transformation of any cell or organism in order to be expressed in this cell or organism and result in an altered, preferably increased oil content of this cell or organism.

15

Figurs

Fig. 1:







Fig 2




SUBSTITUTE SHEET (RULE 26)

Lipid synthesis (pmol/min)



Relative TAG-synthesis



SUBSTITUTE SHEET (RULE 26)

Tables

Tab. 1:

		mol % of added [¹⁴ C]-acyl group in TAG ⁽¹⁾					
Substrate added	[¹⁴ C]-lipid ⁽²⁾	unlabelled lipid ⁽²⁾	1-OH-TAG	2-OH-TAG	1-OH-1-ver-TAG	1-OH-2-ver-TAG	3-OH-TAG
A	mono-[¹⁴ C]-ricinoleoyl-DAG	mono-ricinoleoyl-DAG	2,8	12,4	-	-	-
A	mono-[¹⁴ C]-ricinoleoyl-DAG	mono-vernoleoyl-DAG	3,2	12,1	1,3	-	-
A	mono-[¹⁴ C]-ricinoleoyl-DAG	di-vernoleoyl-DAG	4	10	0,5	1,2	-
A	mono-[¹⁴ C]-ricinoleoyl-DAG	di-ricinoleoyl-PC	0,3	24,8	-	-	-
B	mono-[¹⁴ C]-ricinoleoyl-PC	none	6,8	8,0	-	-	4,7
C	mono-[¹⁴ C]-ricinoleoyl-PC	di-oleoyl-DAG	8,6	9,8	-	-	5,0
C	mono-[¹⁴ C]-ricinoleoyl-PC	mono-ricinoleoyl-DAG	5,7	16,7	-	-	1,9
C	mono-[¹⁴ C]-ricinoleoyl-PC	di-ricinoleoyl-DAG	4,5	9,4	-	-	9,5
C	mono-[¹⁴ C]-ricinoleoyl-PC	mono-vernoleoyl-DAG	6,0	11,5	10,9	0,5	7,4
C	mono-[¹⁴ C]-ricinoleoyl-PC	di-vernoleoyl-DAG	6,7	10,8	1,1	8,4	6,8

Tab. 2:

T1 plant deviation	T2 plant number	nmol fatty acids per mg seed	standard
32-4	1	1277	± 11 (n=2)
	4	1261	± 63 (n=3)
	5	1369	± 17 (n=3)
	6	1312	± 53 (n=4)
	7	1197	± 54 (n=5)
	8	1240	± 78 (n=4)
	9	1283	± 54 (n= 5)
	10	1381	± 35 (n=5)
26-14	1	1444	± 110 (n=4)
	2	1617*	± 109 (n=4)
	3	1374	± 37 (n=2)
	5	1562*	± 70 (n=4)
	6	1393	± 77 (n=4)
	7	1433	± 98 (n=4)
	8	1581*	± 82 (n=4)

Sequence Listing

<210> 1

<211> 1986

<212> genomic DNA

<213> Saccharomyces cerevisiae

<221> CDS

<222> (1)..(1983)

<400> 1

```
atg ggc aca ctg ttt cga aga aat gtc cag aac caa aag agt gat tct      48
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
   1             5             10             15

gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga      96
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
          20             25             30

aac cac att cat cat caa cag gga tta ggc cat aag aga aga agg ggt      144
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
          35             40             45

att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agg      192
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
          50             55             60

aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg      240
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
          65             70             75             80

att ttc att ctt ggt gca ttc tta ggt gta ctt ttg ccg ttt agc ttt      288
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
          85             90             95

ggc gct tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt      336
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
          100            105            110

gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt      384
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
          115            120            125

ctc cca caa ggt ata agt tcg ttt att gat gat att cag gct ggt aac      432
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
          130            135            140
```

tac tcc aca tct tct tta gat gat ctc agt gaa aat ttt gcc gtt ggt	480
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly	
145 150 155 160	
aaa caa ctc tta cgt gat tat aat atc gag gcc aaa cat cct gtt gta	528
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val	
165 170 175	
atg gtt cct ggt gtc att tct acg gga att gaa agc tgg gga gtt att	576
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile	
180 185 190	
gga gac gat gag tgc gat agt tct gcg cat ttt cgt aaa cgg ctg tgg	624
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp	
195 200 205	
gga agt ttt tac atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg	672
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp	
210 215 220	
ttg aaa cat gta atg tta gat cct gaa aca ggt ctg gac cca ccg aac	720
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn	
225 230 235 240	
ttt acg cta cgt gca gca cag ggc ttc gaa tca act gat tat ttc atc	768
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile	
245 250 255	
gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att	816
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile	
260 265 270	
ggc tat gaa ccc aat aaa atg acg agt gct gcg tat gat tgg agg ctt	864
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu	
275 280 285	
gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag	912
Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys	
290 295 300	
gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta	960
Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu	
305 310 315 320	
att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg	1008
Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp	
325 330 335	

SUBSTITUTE SHEET (RULE 26)

gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt	1056
Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val	
340 345 350	
aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc	1104
Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly	
355 360 365	
gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc	1152
Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr	
370 375 380	
att caa tta aat acg tta gcc atg tat ggt ttg gaa aag ttc ttc tca	1200
Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser	
385 390 395 400	
aga att gag aga gta aaa atg tta caa acg tgg ggt ggt ata cca tca	1248
Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser	
405 410 415	
atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct	1296
Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser	
420 425 430	
tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att	1344
Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile	
435 440 445	
cga ttt gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg	1392
Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met	
450 455 460	
aaa gac gcc att aac atg aca tta tcg ata tca cct gaa tgg ctc caa	1440
Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln	
465 470 475 480	
aga aga gta cat gag cag tac tcg ttc ggc tat tcc aag aat gaa gaa	1488
Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu	
485 490 495	
gag tta aga aaa aat gag cta cac cac aag cac tgg tcg aat cca atg	1536
Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met	
500 505 510	
gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac	1584
Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr	
515 520 525	

ggg gtg aac aac cca act gaa agg gca tat gta tat aag gaa gag gat 1632
Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
530 535 540

gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct 1680
Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
545 550 555 560

gta ttc ctc acc gag ggg gac gga acc gtt ccg ctc gtg gcg cat tca 1728
Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
565 570 575

atg tgt cac aaa tgg gcc cag ggt gct tca ccg tac aac cct gcc gga 1776
Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
580 585 590

att aac gtt act att gtg gaa atg aaa cac cag cca gat cga ttt gat 1824
Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
595 600 605

ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctc ggc agc 1872
Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
610 615 620

gcg gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat 1920
Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
625 630 635 640

ctc gtc gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag 1968
Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
645 650 655

atg ccc ttc cca atg taa 1986
Met Pro Phe Pro Met
660

<210> 2

<211> 661

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 2

Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 110
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 140
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 160
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
165 170 175
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180 185 190
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195 200 205
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
210 215 220
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
225 230 235 240
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
245 250 255
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
260 265 270
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
275 280 285
Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
290 295 300
Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
305 310 315 320
Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
325 330 335

SUBSTITUTE SHEET (RULE 26)

Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu
 485 490 495
 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met
 500 505 510
 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr
 515 520 525
 Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
 530 535 540
 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
 545 550 555 560
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
 565 570 575
 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
 580 585 590
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
 595 600 605
 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
 610 615 620
 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
 625 630 635 640
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
 645 650 655
 Met Pro Phe Pro Met
 660

<210> 3
<211> 2312
<212> genomic DNA
<213> Schizosaccharomyces pombe

<400> 3
ATGGCGTCTT CCAAGAAGAG CAAAACATCAT AAGAAAAAGA AAGAAGTCAA 50
ATCTCCTATC GACTTACCAA ATTCAAAGAA ACCAACTCGC GCTTTGAGTG 100
AGCAACCTTC AGCGTCCGAA ACACAATCTG TTTCAAATAA ATCAAGAAAA 150
TCTAAATTTG GAAAAAGATT GAATTTTATA TTGGGCGCTA TTTTGGGAAT 200
ATGCGGTGCT TTTTTTTTCG CTGTTGGAGA CGACAATGCT GTTTTCGACC 250
CTGCTACGTT AGATAAATTT GGGAAATATGC TAGGCTCTTC AGACTTGTTT 300
GATGACATTA AAGGATATTT ATCTTATAAT GTGTTTAAGG ATGCACCTTT 350
TACTACGGAC AAGCCTTCGC AGTCTCCTAG CGGAAATGAA GTTCAAGTTG 400
GTCTTGATAT GTACAATGAG GGATATCGAA GTGACCATCC TGTTATTATG 450
GTTCTGGTG TTATCAGCTC AGGATTAGAA AGTTGGTCGT TTAATAATTG 500
CTCGATTCTT TACTTTAGGA AACGTCTTTG GGGTAGCTGG TCTATGCTGA 550
AGGCAATGTT CCTTGACAAG CAATGCTGGC TTGAACATTT AATGCTTGAT 600
AAAAAAACCG GCTTGGATCC GAAGGGAATT AAGCTGCGAG CAGCTCAGGG 650
GTTTGAAGCA GCTGATTTTT TTATCACGGG CTATTGGATT TGGAGTAAAG 700
TAATTGAAAA CCTTGCTGCA ATTGGTTATG AGCCTAATAA CATGTTAAGT 750
GCTTCTTACG ATTTGGCGGT ATCATATGCA AATTTAGAGG AACGTGATAA 800
ATATTTTTCA AAGTTAAAAA TGTTCAATTGA GTTACAGCAAC ATTGTACATA 850
AGAAAAAGGT AGTGTTGATT TCTCACTCCA TGGGTTTACA GGTACGTAC 900
TATTTTTTTA AGTGGGTTGA AGCTGAGGGC TACGGAAATG GTGGACCGAC 950
TTGGGTTAAT GATCATATTG AAGCATTTAT AAATGTGAGT CTCGATGGTT 1000
GTTTGACTAC GTTTCTAACT TTTGAATAGA TATCGGGATC TTTGATTGGA 1050
GCACCCAAA CAGTGGCAGC GCTTTTATCG GGTGAAATGA AAGATACAGG 1100
TATTGTAATT ACATTAAACA TGTTAATATT TAATTTTTGC TAACCGTTTT 1150
AAGCTCAATT GAATCAGTTT TCGGTCTATG GGTAAGCAAT AAATTGTTGA 1200
GATTTGTTAC TAATTTACTG TTTAGTTTGG AAAAATTTTT TTCCCGTTCT 1250
GAGGTATATT CAAAAATACA AATGTGCTCT ACTTTTTCTA ACTTTTAATA 1300
GAGAGCCATG ATGGTTTCGCA CTATGGGAGG AGTTAGTTCT ATGCTTCCTA 1350
AAGGAGGCGA TGTTGTATGG GGAAATGCCA GTTGGGTAAG AAATATGTGC 1400
TGTTAATTTT TTATTAATAT TTAGGCTCCA GATGATCTTA ATCAAACAAA 1450
TTTTTCCAAT GGTGCAATTA TTCGATATAG AGAAGACATT GATAAGGACC 1500
ACGATGAATT TGACATAGAT GATGCATTAC AATTTTTAAA AAATGTTACA 1550
GATGACGATT TTAAAGTCAT GCTAGCGAAA AATTATTCCC ACGGTCTTGC 1600
TTGGACTGAA AAAGAAGTGT TAAAAAATAA CGAAATGCCG TCTAAATGGA 1650
TAAATCCGCT AGAAGTAAGA ACATTAAAGT TACTAAATTA TACTAACCCA 1700
AATAGACTAG TCTTCCTTAT GCTCCTGATA TGAAAATTTA TTGCGTTCAC 1750
GGGGTCGGAA AACCAACTGA GAGAGGTTAT TATTATACTA ATAATCCTGA 1800
GGGGCAACCT GTCATTGATT CCTCGGTTAA TGATGGAACA AAAGTTGAAA 1850
ATGTGAGAGA ATTTATGTTT CAAACATTCT ATTAAGTGT TTATTAGGGT 1900
ATTGTTATGG ATGATGGTGA TGGAACCTTA CCAATATTAG CCCTTGGTTT 1950
GGTGTGCAAT AAAGTTTGGC AAACAAAAAG GTTTAATCCT GCTAATACAA 2000
GTATCACAAA TTATGAAATC AAGCATGAAC CTGCTGCGTT TGATCTGAGA 2050
GGAGGACCTC GCTCGGCAGA ACACGTCGAT ATACTGGAC ATTCAGAGCT 2100
AAATGTATGT TCATTTTACC TTACAAATTT CTATTACTAA CTCTTGAAAT 2150
AAGGAAATTA TTTTAAAAGT TTCATCAGGC CATGGTGACT CGGTACCAAA 2200
CCGTTATATA TCAGATATCC AGTACGGACA TAAGTTTTGT AGATTGCAAT 2250
TAACTAACTA ACCGAACAGG GAAATAATAA ATGAGATAAA TCTCGATAAA 2300
CCTAGAAATT AA 2312

<210> 4
 <211> 3685
 <212> genomic DNA
 <213> Arabidopsis thaliana

<400> 4

ATGCCCCCTTA	TTCATCGGAA	AAAGCCGACG	GAGAAACCAT	CGACGCCGCC	50
ATCTGAAGAG	GTGGTGCACG	ATGAGGATTC	GCAAAAGAAA	CCACACGAAT	100
CTTCCAAATC	CCACCATAAG	AAATCGAACG	GAGGAGGGAA	GTGGTCGTGC	150
ATCGATTCTT	GTTGTTGGTT	CATTGGGTGT	GTGTGTGTAA	CCTGGTGGTT	200
TCTTCTCTTC	CTTTACAACG	CAATGCCTGC	GAGCTTCCCT	CAGTATGTAA	250
CGGAGCGAAT	CACGGGTCCT	TTGCCTGACC	CGCCCGGTGT	TAAGCTCAAA	300
AAAGAAGGTC	TTAAGGCGAA	ACATCCTGTT	GTCTTCATTC	CTGGGATTGT	350
CACCGGTGGG	CTCGAGCTTT	GGGAAGGCCAA	ACAATGCGCT	GATGGTTTAT	400
TTAGAAAACG	TTTGTGGGGT	GGAACCTTTG	GTGAAGTCTA	CAAAAGGTGA	450
GCTCAACAA	TCTCACTCTT	CCTTTATATT	GGGATTTGGA	TTGGATCTGA	500
TGAGATCACG	CACCTGTTGC	TTCTTCAACA	TCACCTCAAAC	TTTAATTCCA	550
TGTTTGTCTG	TCTTACTCTT	TACTTTTTTTT	TTTTTTTGTAT	GTGAAACGCT	600
ATTTTCTTAA	GAGACTATTT	CTGTATGTGT	AAGGTAAGCG	TTCCAAGGAC	650
GTAATTGGCT	TGGACTATTT	CTGTTTGATT	GTAACTTTA	GGATATAAAA	700
TAGCTGCCTT	GGAATTTCAA	GTCATCTTAT	TGCCAAATCT	GTGCTAGAC	750
ATGCCCTAGA	GTCCGTTTCA	AACAAGTTAC	TTCTTTTACT	GTCTTGCCT	800
GTAGATTTAG	CTTGTGTAG	CGTATAATGA	AGTAGTGTTT	TATGTTTGT	850
TGGGAATAGA	GAAGTTCTAA	CTACATCTGT	GGAAAGTGTG	TTCAGGCTGT	900
GATAGAGGAC	TGTTGCTTTA	TTATTCAACT	ATGTATATGT	GTAATTAAAG	950
CTAGTTCCTT	TTTGATCTTT	CAGCTCAATG	TGCTTTTCTC	AATTTTTTTC	1000
TCAATTTCAA	AGTTTCACAT	CGAGTTTATT	CACATGTCTT	GAATTTTCGT	1050
CATCCTCGTT	CTGTTATCCA	GCTTTGAACT	CCTCCCGACC	CTGCTATGGA	1100
TATATTAAAA	AAAAAGTGTT	TTGTGGGTTG	CATCTTTGTT	ACGATCTGCA	1150
TCTTCTTCTT	TCGGCTCAGT	GTCATGTTT	TTGCTATGGT	AGAGATGGGC	1200
AATGTTATTG	TTGATGGTAA	CAGTGGTATA	GTTGATAGTA	TCTTAACATA	1250
TCAATTATCT	CTTTGATTCA	GGCCTCTATG	TTGGGTGGAA	CACATGTCAC	1300
TTGACAATGA	AACCTGGGTTG	GATCCAGCTG	GTATTAGAGT	TCGAGCTGTA	1350
TCAGGACTCG	TGGCTGCTGA	CTACTTTGCT	CCTGGCTACT	TTGTCTGGGC	1400
AGTGCTGATT	GCTAACCTTG	CACATATTGG	ATATGAAGAG	AAAAATATGT	1450
ACATGGCTGC	ATATGACTGG	CGGCTTTCTG	TTCAGAACAC	AGAGGTTCTT	1500
TTCTCATCGT	TCTTTCTATT	ATTCTGTTCC	ATGTTACGTT	TCTTTCTTCA	1550
TTACTTAAGG	CTTAAATATG	TTTCATGTTG	AATTAATAGG	TACGTGATCA	1600
GACTCTTAGC	CGTATGAAAA	GTAATATAGA	GTGATGGTT	TCTACCAACG	1650
GTGGAAAAAA	AGCAGTTATA	GTTCCGCATT	CCATGGGGGT	CTTGATTTTT	1700
CTACATTTTA	TGAAGTGGGT	TGAGGCACCA	GCTCCTCTGG	GTGGCGGGGG	1750
TGGGCCAGAT	TGGTGTGCAA	AGTATATTAA	GGCGGTGATG	AACATTGGTG	1800
GACCATTTCT	TGGTGTGTTA	AAAGCTGTTG	CAGGGCTTTT	CTCTGCTGAA	1850
GCAAAGGATG	TTGCTAGTTG	CAGGTATTGA	ATATCTGCTT	ATACTTTTGA	1900
TGATCAGAAC	CTTGCTCTG	GAACCTCAAAG	TTATTCTACT	AAATATCAAT	1950
TCTAATAACA	TTGCTATATT	ATCGCTGCAA	CTGACATTGG	TTGATTATTT	2000
TTGCTGCTTA	TGTAACGAA	ACTCTCTTGA	GATTAGACAA	ATGATGAATT	2050
GATAATTCTT	ACGCATTGCT	CTGTGATGAC	CAGTTTCTTA	GCTTCGACGA	2100
TAACATTTGT	CATACTGTCT	TTTGGAGGGC	ATTGAATTTT	GCTATGGAAA	2150
GCGCTGGAGC	TTCCATGCTT	GCATTCTTTA	CCAATTAGCG	TTATTCTGCT	2200
TCTTTCAATT	TTCTTGATA	TGCATCTATG	GTCTTTTATT	TCTTCTTAAT	2250
TAAAGACTCG	TTGGATTAGT	TGCTCTATTA	GTCATTGGT	TCCTTAATAT	2300
AGAACTTTAC	TTTCTTCGAA	AATTGCAGAG	CGATTGCCCC	AGGATTCTTA	2350
GACACCGATA	TATTTAGACT	TCAGACCTTG	CAGCATGTAA	TGAGAATGAC	2400
ACGCACATGG	GACTCAACAA	TGTCTATGTT	ACCGAAGGGA	GGTGACACGA	2450
TATGGGGCGG	GCTTGATTGG	TCACCGGAGA	AAGGCCACAC	CTGTTGTGGG	2500
AAAAAGCAAA	AGAACAACGA	AACTTGTGGT	GAAGCAGGTG	AAAACGGAGT	2550
TTCCAAGAAA	AGTCCTGTTA	ACTATGGAAG	GATGATATCT	TTTGGGAAAG	2600
AAGTAGCAGA	GGCTGCGCCA	TCTGAGATTA	ATAATATTGA	TTTTCGAGTA	2650
AGGACATATA	AATCATAATA	AACCTTGTTT	ATTTTGTGAT	TGTATGATGA	2700
ATATCTGTAC	ATTTTATCTG	GTGAAGGGTG	CTGTCAAAGG	TCAGAGTATC	2750
CCAAATCACA	CCTGTCGTGA	CGTGTGGACA	GAGTACCATG	ACATGGGAAT	2800
TGCTGGGATC	AAAGCTATCG	CTGAGTATAA	GGTCTACACT	GCTGGTGAAG	2850

SUBSTITUTE SHEET (RULE 26)

CTATAGATCT	ACTACATTAT	GTTGCTCCTA	AGATGATGGC	GCGTGGTGCC	2900
GCTCATTTCT	CTTATGGAAT	TGCTGATGAT	TTGGATGACA	CCAAGTATCA	2950
AGATCCCAAA	TACTGGTCAA	ATCCGTTAGA	GACAAAGTAA	GTGATTTCTT	3000
GATTCCAACT	GTATCCTTCG	TCCTGATGCA	TTATCAGTCT	TTTTGTTTTT	3050
GGTCTTGTG	GATATGGTTT	TCAGCTCAAA	GCTTACAAAG	CTGTTTCTGA	3100
GCCTTTCTCA	AAAAGGCTTG	CTCAGTAATA	TTGAGGTGCT	AAAGTTGATA	3150
CATGTGACTC	TTGCTTATAA	ATCCTCCGTT	TGGTTTGTTT	TGCTTTTTTCA	3200
GATTACCGAA	TGCTCCTGAG	ATGGAAATCT	ACTCATTATA	CGGAGTGGGG	3250
ATACCAACGG	AACGAGCATA	CGTATACAAG	CTTAACCAGT	CTCCCGACAG	3300
TTGCATCCCC	TTTCAGATAT	TCACTTCTGC	TCACGAGGAG	GACGAAGATA	3350
GCTGTCTGAA	AGCAGGAGTT	TACAATGTGG	ATGGGGATGA	AACAGTACCC	3400
GTCTTAAGTG	CCGGGTACAT	GTGTGCAAAA	GCGTGGCGTG	GCAAGACAAG	3450
ATTCAACCCCT	TCCGGAATCA	AGACTTATAT	AAGAGAATAC	AATCACTCTC	3500
CGCCGGCTAA	CCTGTTGGAA	GGGCGCGGGA	CGCAGAGTGG	TGCCCCATGTT	3550
GATATCATGG	GAAACTTTGC	TTTGATCGAA	GATATCATGA	GGGTTGCCGC	3600
CGGAGGTAAC	GGGTCTGATA	TAGGACATGA	CCAGGTCCAC	TCTGGCATAT	3650
TTGAATGGTC	GGAGCGTATT	GACCTGAAGC	TGTGA		3685

<210> 5
<211> 2427
<212> cDNA
<213> Arabidopsis thaliana

<400> 5
AGAAACAGCT CTTTGTCTCT CTCGACTGAT CTAACAATCC CTAATCTGTG 50
TTCTAAATTC CTGGACGAGA TTGACAAAG TCCGTATAGC TTAACCTGGT 100
TTAATTTCAA GTGACAGATA TGCCCTTAT TCATCGGAAA AAGCCGACGG 150
AGAAACCATC GACGCCGCCA TCTGAAGAGG TGGTGCACGA TGAGGATTCG 200
CAAAAGAAAC CACACGAATC TTCCAAATCC CACCATAAGA AATCGAACGG 250
AGGAGGGAAG TGGTCGTGCA TCGATTCTTG TTGTTGGTTC ATTGGGTGTG 300
TGTGTGTAAC CTGGTGGTTT CTTCTCTTCC TTTACAACGC AATGCCTGCG 350
AGCTTCCCTC AGTATGTAAC GGAGCGAATC ACGGGTCCTT TGCCTGACCC 400
GCCCCGGTGT AAGCTCAAAA AAAGAAGGTC TTAAGGCGAA ACATCCTGTT 450
GTCCTTCATTC CTGGGATGTG CACCGGTGGG CTCGAGCTTT GGAAGGCAA 500
ACAATGCGCT CATGGTTTAT TTAGAAAACG TTTGTGGGGT GGAACTTTGT 550
GTGAAGTCTA GAAAAGGCCT CTATGTTGGG TGGAAACACAT GTCACCTGAC 600
AATGAAACTG GGTGGGATCC AGCTGGTATT AGAGTTCGAG CTGTATCAGG 650
ACTCGTGGCT GCTGACTACT TTGCTCCTGG CTACTTTGTC TGGGCAGTGC 700
TGATTGCTAA CCTTGCACAT ATTGGATATG AAGAGAAAAA TATGTACATG 750
GCTGCATATG ACTGGCGGCT TTCGTTTCAG AACACAGAGG TACGTGATCA 800
GACTCTTAGC CGTATGAAAA GTAATATAGA GTTGATGGTT TCTACCAACG 850
GTGGAAAAAA AGCAGTTATA GTTCCGCATT CCATGGGGGT CTTGTATTTT 900
CTACATTTTA TGAAGTGGGT TGAGGCACCA GTCCTCTGCG GTGGCGGGGG 950
TGCGCCAGAT TGGTGTGCAA AGTATATTAA GGCGGTGATG AACATTGGTG 1000
GACCATTCTT TTGCTTTCCA AAAGCTGTTG CAGGGCTTTT CTCTGCTGAA 1050
GCAAAGGATG TTGTCAGTTGC CAGAGCGATT GCCCCAGGAT TCTTAGACAC 1100
CGATATATTT AGACTTCAGA CCTTGCAGCA TGTAATGAGA ATGACACGCA 1150
CATGGGACTC AACAATGTCT ATGTTACCGA AGGGAGGTGA CACGATATGG 1200
GGCGGGCTTG ATTGGTCACC GGAGAAAGGC CACACCTGTT GTGGGAAAAA 1250
GCAAAAGAAC AACGAAACTT GTGGTGAAGC AGGTGAAAAC GGAGTTTCCA 1300
AGAAAAGTCC TGTTAACTAT GGAAGGATGA TATCTTTTGG GAAAGAAGTA 1350
GCAGAGGCTG CGCCATCTGA GATTAATAAT ATTGATTTTC GAGGTGCTGT 1400
CAAAGGTACAG AGTATCCCAA ATCACACCTG TCGTGACGTG TGGACAGAGT 1450
ACCATGACAT GGAATTGCT GGGATCAAAG CTATCGCTGA GTATAAGGTC 1500
TACACTGCTG GTGAAGCTAT AGATCTACTA CATTATGTTG CTCCTAAGAT 1550
GATGGCGCGT GGTGCCGCTC ATTTCTCTTA TGGAATTGCT GATGATTTGG 1600
ATGACACCAA GTATCAAGAT CCCAAATACT GGTCAAATCC GTTAGAGACA 1650
AAATTACCGA ATGCTCCTGA GATGGAAATC TACTCATTAT ACGGAGTGGG 1700
GATACCAACG GAACGAGCAT ACGTATACAA GCTTAACCAG TCTCCCGACA 1750
GTTGCATCCC CTTTCAGATA TTCACTTCTG CTCACGAGGA GGACGAAGAT 1800
AGCTGTCTGA AAGCAGGAGT TTACAAATGT GATGGGGATG AAACAGTACC 1850
CGTCCTAAGT GCCGGGTACA TGTGTGCAAA AGCGTGCGGT GGCAAGACAA 1900
GATTCACACC TTCCGGAATC AAGACTTATA TAAGAGAATA CAATCACTCT 1950
CCGCCGGCTA ACCTGTTTGA AGGGCGCGGG ACGCAGAGTG GTGCCCATGT 2000
TGATATCATG GGAAACTTTG CTTTGATCGA AGATATCATG AGGGTTGCCG 2050
CCGGAGGTAA CGGGTCTGAT ATAGGACATG ACCAGGTCCA CTCTGGCATA 2100
TTTGAATGGT CGGAGCGTAT TGACCTGAAG CTGTGAATAT CATGATCTCT 2150
TTAAGCTGTC CTGTCAGCTT ATGTGAATCC AATACTTTGA AAGAGAGATC 2200
ATCATCAATT CATCATCATC GTCATCATCA TGATGCTCAA CTCACAAAGA 2250
AGCCTGAGAA TGATACTTTG GTGCGAAATT CTCAATACCT CTTTAATATT 2300
CTTATTGAAT GTAAATTATA CAATCCTATC TAATGTTTGA ACGATAACAC 2350
AAAACCTGCT GCNGCCATGT TTGTTGTCT TGTCAAAGC ATCAATTGT 2400
GGGTAAAAA AAAAAAAAAA AAAAAA 2427

SUBSTITUTE SHEET (RULE 26)

<210> 6
<211> 671
<212> PRT
<213> Arabidopsis thaliana

<400> 6
MPLIHRKKPT EKPSTPPSEE VVHDEDSQKK PHESSKSHHK KSNGGGKWSC 50
IDSCCWFFIGC VCVTWWFLLF LYNAMPASFP QYVTERITGP LPDPPGVKLK 100
KEGLKAKHPV VFIPGIVTGG LELWEGKQCA DGLFRKRLWG GTFGEVYKRP 150
LCWVEHMSLD NETGLDPAGI RVRVAVSGLVA ADYFAPGYFV WAVLIANLAH 200
IGYEENMYM AAYDWRLSFQ NTEVRDQTLS RMKSNIELMV STNGGKKAVI 250
VPHSMGVLYF LHFMKWVEAP APLGGGGGPD WCAKYIKAVM NIGGPFLGVP 300
KAVAGLFSAE AKDVAVARAI APGFLDTDIF RLQTLQHVMR MTRTWDSTMS 350
MLPKGGDTIW GGLDWSPEKG HTCCGKKQKN NETCGEAGEN GVSCKSPVNY 400
GRMISFGKEV AEAAPSEINN IDFRGAVKGQ SIPNHTCRDV WTEYHDMGIA 450
GIKAI AEYKV YTAGEAIDLL HYVAPKMMAR GAAHFSYGIA DDLDDTKYQD 500
PKYWSNPLET KLPNAPEMEI YSLYGVGIPT ERAYVYKLNQ SPDSCIPFQI 550
FTSAHEEDED SCLKAGVYNV DGDETVFVLS AGYMCAKAWR GKTRFNPSGI 600
KTYIREYNHS PPANLLEGRG TQSGAHVDIM GNFALIEDIM RVAAGGNGSD 650
IGHDQVHSGI FEWSERIDLK L 671

<210> 7

<211> 643

<212> cDNA

<213> Zea mays

<221> CDS

<222> (1)..(402)

<400> 6

CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT 48

Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly

1

5

10

15

GAT GAA ACT GTT CCA GTT CTT AGT GCG GGC TAC ATG TGT GCG AAA GGA 96

Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly

20

25

30

TGG CGT GGC AAA ACT CGT TTC AGC CCT GCC GGC AGC AAG ACT TAC GTG 144

Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val

35

40

45

AGA GAA TAC AGC CAT TCG CCA CCC TCT ACT CTC CTG GAA GGC AGG GGC 192

Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly

50

55

60

ACC CAG AGC GGT GCA CAT GTT GAT ATA ATG GGG AAC TTT GCT CTA ATT 240

Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile

65

70

75

80

GAG GAC GTC ATC AGA ATA GCT GCT GGG GCA ACC GGT GAG GAA ATT GGT 288

Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly

85

90

95

GGC GAT CAG GTT TAT TCA GAT ATA TTC AAG TGG TCA GAG AAA ATC AAA 336

Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys

100

105

110

TTG AAA TTG TAA CCTATGGGAA GTTAAAGAAG TGCCGACCCG TTTATTGCGTTCC 391

Leu Lys Leu

115

AAAGTGTCCT GCCTGAGTGC AACTCTGGAT TTTGCTTAAA TATTGTAATT TTTCACGC 449

TTCATTCGTC CCTTTGTCAA ATTTACATTT GACAGGACGC CAATGCGATA CGATGTTG 507

TACCGCTATT TTCAGCATTG TATATTAAAC TGTACAGGTG TAAGTTGCAT TTGCCAGC 565

TGAAATTGTG TAGTCGTTTT CTTTACGATT TAATANCAAG TGGCGGAGCA GTGCCCCA 623

AGCNAAAAAA AAAAAAAAAA

643

<210> 8

<211> 115

<212> PRT

<213> Zea mays

<400> 8

Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
1 5 10 15

Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
20 25 30

Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
35 40 45

Arg Glu Tyr Ser His Ser Pro Ser Thr Leu Leu Glu Gly Arg Gly
50 55 60

Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
65 70 75 80

Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
85 90 95

Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
100 105 110

Leu Lys Leu
115

<210> 9

<211> 616

<212> cDNA

<213> Neurospora crassa

<400> 9

ggtggcgaag	acganggcgg	aagttggagg	ctaacgagaa	tgacnctcgg	50
agatggatct	accctctaga	gacacgacta	ccnttgcacc	cagcctcaag	100
gtntacngtt	tntatgggta	ggaagccgac	ggagcgagcc	tacatctatc	150
tggcgcccga	tcccgggacg	acaacgcac	tttagatgac	gacgatacg	200
actttgactn	aggggcacat	tgaccacggt	gtgatttttg	gcgaaggcga	250
tggcacagtg	aaccttatga	gtttggggta	cctgtgcaat	aaggggtgga	300
aaatgaagag	atacaatcct	gcgggctcaa	aaataaccgt	ggcgcgagatg	350
ccgcatgaac	cagaacgggt	caatccgaga	ggagggccga	atacggcgga	400
tcacgtggat	attctaggaa	ggcagaatct	aaacgagtac	attcttaaag	450
tggcggcagg	tcgaggcgat	acaattgagg	attttattac	tagtaatatt	500
cttaaatatg	tagaaaaggt	tgaaatttat	gaagagtaat	taaatacggc	550
acataggtta	ctcaatagta	tgactaatta	aaaaaaaaatt	ttttttctaa	600
aaaaaaaaaa	aaaaaa				616

<210> 10
<211> 1562
<212> genomic DNA
<213> Arabidopsis thaliana

<400> 10
ATGAAAAAAA TATCTTCACA TTATTCGGTA GTCATAGCGA TACTCGTTGT 50
GGTGACGATG ACCTCGATGT GTCAAGCTGT GGGTAGCAAC GTGTACCCTT 100
TGATTCTGGT TCCAGGAAAC GGAGGTAACC AGCTAGAGGT ACGGCTGGAC 150
AGAGAATACA AGCCAAGTAG TGTCTGGTGT AGCAGCTGGT TATATCCGAT 200
TCATAAGAAG AGTGGTGGAT GGTTTAGGCT ATGGTTTCGAT GCAGCAGTGT 250
TATTGTCTCC CTTCAACAGG TGCTTCAGCG ATCGAATGAT GTTGTACTAT 300
GACCCTGATT TGGATGATTA CCAAAATGCT CCTGGTGTCC AAACCCGGGT 350
TCCTCATTTT GGTTCGACCA AATCACTTCT ATACCTCGAC CCTCGTCTCC 400
GGTTAGTACT TTCCAAGATA TATCATTTTG GGACATTTGC ATAATGAACA 450
AAATAGACAT AAATTTGGGG GATTATTGTT ATATCAATAT CCATTTATAT 500
GCTAGTCGGT AATGTGAGTG TTATGTTAGT ATAGTTAATG TGAGTGTTAT 550
GTGATTTTCC ATTTTAAATG AAGCTAGAAA GTTGTTCGTTT AATAATGTTG 600
CTATGCTATG AGAATTATAA GGACACTATG TAAATGTAGC TTAATAATAA 650
GGTTTGATTT GCAGAGATGC CACATCTTAC ATGGAACATT TGGTGAAAGC 700
TCTAGAGAAA AAATGCGGGT ATGTTAACGA CCAAACCATC CTAGGAGCTC 750
CATATGATTT CAGGTACGGC CTGGCTGCTT CGGGCCACCC GTCCCGTGTA 800
GCCTCACAGT TCCTACAAGA CCTCAAACAA TTGGTGGAAA AAACCTAGCAG 850
CGAGAACGAA GGAAAGCCAG TGATACTCCT CTCCCATAGC CTAGGAGGAC 900
TTTTCGTCCT CCATTTCTCT AACCCTACCA CCCCTTCATG GCGCCGCAAG 950
TACATCAAAC ACTTTGTTGC ACTCGCTGCG CCATGGGGTG GGACGATCTC 1000
TCAGATGAAG ACATTTGCTT CTGGCAACAC ACTCGGTGTC CCTTTAGTTA 1050
ACCCCTTGCT GGTCAGACGG CATCAGAGGA CCTCCGAGAG TAACCAATGG 1100
CTACTTCCAT CTACCAAAAGT GTTTCACGAC AGAACTAAAC CGCTTGTCGT 1150
AACTCCCCAG GTTAACTACA CAGCTTACGA GATGGATCGG TTTTTTGCAG 1200
ACATTGGATT CTCACAAGGA GTTGTGCCTT ACAAGACAAG AGTGTTCCTT 1250
TTAACAGAGG AGCTGATGAC TCCGGGAGTG CCAGTCACTT GCATATATGG 1300
GAGAGGAGTT GATACACCGG AGGTTTTGAT GTATGGAAA GGAGGATTCG 1350
ATAAGCAACC AGAGATTAAG TATGGAGATG GAGATGGGAC GGTTAATTTG 1400
GCGAGCTTAG CAGCTTTGAA AGTCGATAGC TTGAACACCG TAGAGATTGA 1450
TGGAGTTTCC CATACTCTA TACTTAAAGA CGAGATCGCA CTTAAAGAGA 1500
TTATGAAGCA GATTTCAATT ATTAATTATG AATTAGCCAA TGTTAATGCC 1550
GTCAATGAAT GA 1562

<210> 11
<211> 3896
<212> genomic DNA
<213> Arabidopsis thaliana

<400> 11

ATGGGAGCGA	ATTCGAAATC	AGTAACGGCT	TCCTTCACCG	TCATCGCCGT	50
TTTTTTCTTG	ATTTGCGGTG	GCCGAACGTC	GGTGGAGGAT	GAGACCGAGT	100
TTCACGGCGA	CTACTCGAAG	CTATCGGGTA	TAATCATTCC	GGGATTTGCG	150
TCGACGCAGC	TACGAGCGTG	GTCGATCCTT	GACTGTCCAT	ACACTCCGTT	200
GGACTTCAAT	CCGCTCGACC	TCGTATGGCT	AGACACCACT	AAGGTCCGTG	250
ATCTTCATTT	CCTTCGCTCC	TTATTCTGTC	GGTCGAGTCA	CTTGTTGATG	300
AATTCCAAGC	GAAATATAGC	AATGAAGCAT	GTCTCGTCTC	TCTTATTGAT	350
TCGTTCAATTA	GTCAACAGTG	ACGCTTCTGA	ATCTGAGTTT	AGAGTCATAT	400
AAAACAGCTG	ACTCGGCGAG	TGTTTCCCAT	CGCTTTTGGT	TCGCTAAATG	450
TAGCGCAATG	AATGTGTAAT	TAGTCTGCGC	TTTTTATTCA	ACTAGATCTG	500
CAAGTTTTC	AGAGTGCTCA	ATAGTAGTTA	GAAAATGTGA	GGTCATTTTA	550
CTTGTCGATT	GTGATTCTTT	TGGTTGTTGC	TTACTGATCG	ACGTGATGGA	600
TGGTTTACAG	CTTCTTTCTG	CTGTCAACTG	CTGGTTTAAG	TGTATGGTGC	650
TAGATCCTTA	TAATCAAACA	GACCATCCCG	AGTGTAAGTC	ACGGCCTGAC	700
AGTGGTCTTT	CAGCCATCAC	AGAATTGGAT	CCAGGTTACA	TAACAGGTAG	750
TTTCGGATTT	TTCTTTCTTT	TGAGTTTCT	TCAATTTGAT	ATCATCTTGT	800
TGTGATATAA	TATGGCTAAG	TTCATTAATT	TGGTCAATTT	TCAGGTCCCTC	850
TTTCTACTGT	CTGGAAAGAG	TGGCTTAAGT	GGTGTGTGA	GTTTGGGTATA	900
GAAGCAAATG	CAATTGTGCG	TGTTCCATAC	GATTGGAGAT	TGTCACCAAC	950
CAAATTTGGA	GAGCGTGACC	TTTACTTTCA	CAAGCTCAAG	TTAGTCCTTA	1000
TCAGGCTAAT	GTCTTTTATC	TTCTCTTTTT	ATGTAAGATA	AGCTAAGAGC	1050
TCTGGTCGTC	TTCTTTTTTG	CAGGTTGACC	TTTGAAACTG	CTTTAAAACT	1100
CCGTGGCGGC	CCTTCTATAG	TATTTGCCCA	TTCAATGGGT	AATAATGTCT	1150
TCAGATACTT	TCTGGAATGG	CTGAGGCTAG	AAATTGCACC	AAAACATTAT	1200
TTGAAGTGGC	TTGATCAGCA	TATCCATGCT	TATTTGCGCTG	TTGGTACCGG	1250
CCTACTATCC	TTAAGTTACC	ATTTTATTTT	TTCTCTAATT	GGGGGAGTTA	1300
TGTTGTGACT	TACTGGATTG	AGCTCGATAC	CTGATTTGTT	GTTGATTTAG	1350
GAGCTCCTCT	TCTTGGTTCT	GTTGAGGCAA	TCAAATCTAC	TCTCTCTGGT	1400
GTAACGTTTG	GCCTTCTGT	TTCTGAGGTG	ACCTCTGACT	TCTCTTTAGT	1450
TTTAAGTAGT	TGATATCAAC	CAGGTCTTAT	AACTCACTGG	ATTTTCTTTT	1500
TGAAAGTATT	ACTTTTGTTA	ATTGAACTGC	TGTACGCGAT	ATGGTATCTG	1550
TAGATCTTGA	AGTGCTAGTT	ATCAAAGAAC	ATATTGTGGG	TAGTATACCT	1600
GTCAGCGGCC	TTAGCTAATA	CAACCAAACC	ACATGTACAC	TGATTTAGTT	1650
TTCAGATTAT	TATGGTAGAC	TTTAAGTTGA	GAAGAAACTT	TGACTGAAAT	1700
CTTTTATTTT	TAATAGGCTA	TGATTTGTTT	ATTGAAATCA	TGTGACATAT	1750
TGACATGCGC	TTCTCATGTT	TTTGTGTGGC	AAGGCTTCAG	GGAAGTGCTC	1800
GGTTGTTGTC	CAATTCTTTT	GCGTCGTCAT	TGTGGCTTAT	GCCATTTTCA	1850
AAGAATTGCA	AGGGTGATAA	CACATTCTGG	ACGCATTTTT	CTGGGGGTGC	1900
TGCAAAGAAA	GATAAGCGCG	TATACCACTG	TGATGAAGAG	GAATATCAAT	1950
CAAAATATTC	TGGCTGGCCG	ACAAATATTA	TTAACATTGA	AATTCCTTCC	2000
ACTAGCGGTT	AGACTCTGTA	TATGCAACTG	TAACACTAAC	AAAAGTTTCA	2050
CCAAGAATGT	TCACTCTCAT	ATTTCTGTTCC	TTTGATGTGT	ATCCATCAGT	2100
TACAGAAACA	GCTCTAGTCA	ACATGACCAG	CATGGAATGT	GGCCTTCCCA	2150
CCCTTTTGTC	TTTCACAGCC	CGTGAAC TAG	CAGATGGGAC	TCTTTTCAAA	2200
GCAATAGAAG	ACTATGACCC	AGATAGCAAG	AGGATGTTAC	ACCAGTTAAA	2250
GAAGTACGTA	CCTTCTTTTG	TGATAAGAAA	TATTGCTCAT	CGATCATCAC	2300
TTGCTGGCTT	CTTGACGCT	AAATTGTTTT	GTTTAAATCT	CTATATCAAT	2350
TGTTCAATAG	CCTTGCTTTT	CTTACTATAA	GAAACAAGTA	TAATCAGAAA	2400
CCTTATTATT	GATTATCAGT	TCTCTCCTTA	TATTATGGAA	TGTCTTTTTC	2450
GTTTACAGTT	ATGAATGCAA	AAGGGGGTAT	TTTAGTTGAT	TGATTCTCTC	2500
ATTCTCTAGT	TTGTTTTGAC	TAATAGCGTC	AATTTTGTTT	TTCTAGCAAA	2550
TCTTTGTGAA	TTATATATAA	CATGCTAACT	ATACTTTTCA	GGTTGTATCA	2600
TGATGACCCT	GTTTTTAATC	CTCTGACTCC	TTGGGAGAGA	CCACCTATAA	2650
AAAATGTATT	TTGCATATAT	GGTGCTCATC	TAAAGACAGA	GGTATGATGC	2700
ATTCTCAATA	TCACATTATG	CGTTGACTTT	GTTATTATAT	TCCCCATTTG	2750
GTTTGCAATA	TCTTTTGTAA	TTATGATTTA	TCTTCTCCCT	TGCATCTTAT	2800
GCTATTAAAGC	GTTAAAGGTA	CTAAATGTAT	GAAGCTGTCT	GTCATAGGTT	2850

SUBSTITUTE SHEET (RULE 26)

GGTTATTACT	TTGCCCCAAG	TGGCAAACCT	TATCCTGATA	ATTGGATCAT	2900
CACGGATATC	ATTTATGAAA	CTGAAGGTTC	CCTCGTGTCA	AGGTAATTTT	2950
CCGCAATGGC	AGAAGTAAAA	CAGGAAGGCA	AAGTCTTCTG	TATCAGTCTA	3000
GTGGCATGTT	ATCTCAGTTG	CATAAGCAAA	TTATTAAACA	ACTAAAATTT	3050
AAGTACTTTT	TTATCATTCC	TTTTGAGCTT	AGTGGATGAT	CAGTGGCTTA	3100
AAGTGGGAAG	AGGTGTTGCA	TGAAACATGA	CACTTGATC	AAAGATAACT	3150
AGCAAAACAA	AACTAACCCA	TTTCTGAATT	TCATATTATT	AGGAGTAGTC	3200
GTGCTTTTAA	AAAATTTGTT	TTAAGAAACC	GAAAAACTAG	TTCATATCTT	3250
GATTGTGCAA	TATCTGCAGG	TCTGGAACCTG	TGGTTGATGG	GAACGCTGGA	3300
CCTATAACTG	GGGATGAGAC	GGTAAGCTCA	GAAGTTGGTT	TTGAAATTAT	3350
CTTCTTGCAA	ACTACTGAAG	ACTAAGATAA	TACTTGCTTC	TGGAACACTG	3400
CTTGCTATGT	TCTCTAGTAC	ACTGCAATAT	TGACTCTCCG	CTACTTTTAT	3450
TGATTATGAA	ATTGATCTCT	TATAGGTACC	CTATCATTCA	CTCTCTTGGT	3500
GCAAGAATTG	GCTCGGACCT	AAAGTTAACA	TAACAATGGC	TCCCCAGGTA	3550
CTCTTTTTTA	GTTCTCACC	TTATATAGAT	CAAACTTTAA	GTGTACTTTT	3600
CTGGTTATGT	GTTGATTTAC	CTCCAATTTG	TTCTTTCTAA	AAATCATATA	3650
TCTCTGTACT	CCTCAAGAAC	TTGTATTAAT	CTAAACGAGA	TTCTCATTGG	3700
GAAAATAAAA	CAACAGCCAG	AACACGATGG	AAGCGACGTA	CATGTGGAAC	3750
TAAATGTTGA	TCATGAGCAT	GGGTCAGACA	TCATAGCTAA	CATGACAAAA	3800
GCACCAAGGG	TTAAGTACAT	AACCTTTTAT	GAAGACTCTG	AGAGCATTCC	3850
GGGGAAGAGA	ACCGCAGTCT	GGGAGCTTGA	TAAAAGTGGG	TATTAA	3896

<210> 12
<211> 709
<212> cDNA
<213> Lycopersicon esculentum

<400> 12
CTGGGGCCAA AAGTGAACAT AACAAGGACA CCACAGTCAG AGCATGATGT 50
TCAGATGTAC AAGTGCATCT AAATATAGAG CATCAACATG GTGAAGATAT 100
CATTCCCAAT ATGACAAAGT TACCTACAAT GAAGTACATA ACCTATTATG 150
AGGATTCTGA AAGTTTTCCA GGGACAAGAA CAGCAGTTTG GGAGCTTGAT 200
AAAGCAAATC ACAGGAACAT TGTGAGATCT CCAGCTTTGA TGCGGGAGCT 250
GTGGCTTGAG ATGTGGCATG ATATTCATCC TGATAAAAAG TCCAAGTTTG 300
TTACAAAAGG TGGTGTCTGA TCCTCACTAT TTTCTTCTAT AAATGTTTGA 350
GTTTGTATTG ACATTGTAAG TATTGCAACA AAAAGCAAAG CGTGGGCCTC 400
TGAGGGATGA GGACTGCTAT TGGGATTACG GGAAAGCTCG ATGTGCATGG 450
GCTGAACATT GTGAATACAG GTTAGAATAT TCAAATTATA TTTTGCAAAA 500
TATTCTCTTT TTGTGTATTT AGGCCACCTT TCCCCGGTCA CAACGATGCA 550
GATATGTATT CGGGGATGTT CACCTGGGAC AGAGTTGCAG ATTGAAGAGT 600
TCTACATCTC ACATCCTGTC ACACTATGTG TGATATTTAA GAAACTTTGT 650
TTGGCGGAAC AACAAGTTTG CACAAACATT TGAAGAAGAA AGCGAAATGA 700
TTCAGAGAG 709

<210> 13
<211> 623
<212> PRT
<213> *Schizosaccharomyces pombe*

<400> 13

MASSKKS KTHKKKKEVKSPIDL PNSKKPTRALSEQPSASETQSVSNKSRKSKFGKRLNFILGAILGICGA70
FFFAVGDDNAVFD PATLDKFGNMLGSSDLFDDIKGYLSYNVFKDAPFTTDKPSQSPSGNEVQVGLDMYNE140
GYRSDHPVIMVPGVISSGLESWSFNCSIPYFRKRLWGSWSMLKAMFLDKQCWLEHMLDKKTGLDPKGI210
KLRAAQGF EAADFFITGYWIWSKV IENLAAIGYEPNNMLSASYDWRLSYANLEERDKYFSKLKMFIEYSN280
IVHKKKVVLISHSMGSQVTTYFFKWVEAEGYGNGGPTWVNDHIEAFINISGSLIGAPKTVAALLSGEMKD350
TGIVITL NILEKFFSRSERAMMVRTMGGVSSMLPKGGDVAPDDL NQTNFSNGAIIRYREDIDKD HDEFDI420
DDALQFLKNVTDDDFKVMLAKNYS HGLAWTEKEVLKNNEMPSKWINPLETSLPYAPDMKIYCVHGVGKPT490
ERGYYYTNNPEGQPVIDSSVNDG TKVENGI VMDDGDGTLPI LALGLVCNKVWQTKRFNPANTSITNYEIK560
HEPAAFDLRGGPRSAEHVDILGHSELNEIILKVSSGHGDSVPNRYISDIQEINEINLDKPRN 623

<210> 14

<211> 432

<212> PRT

<213> *Arabidopsis thaliana*

<400> 14

MKKISSHYSVVIAILVVVTMTSMCQAVGSNVYPLILVPGNGGNQLEVRDREYKPSSVWCSSWLYPEIHKK70
SGGWFRWLWFDAAVLLSPFTRCFSDRMMLYYDPDLDDYQNAFGVQTRVPHFGSTKSLLYLDPRLRDATSYM140
EHLVKALEKKCGYVNDQITLGAPYDFRYGLAASGHPSRVASQFLQDLKQLVEKTSSNEGKPVILLSHSL210
GGLFVLHFLNRTTPSWRRKYIKHFVALAAPWGGTISQMKTFASGNTLGVPVLVNPLLVRRHQRTSESQWL280
LPSTKVFHDKPLVVTPOVNYTAYEMDRFFADIGFSQGVVPYKTRVLPLTEELMTPGVPVTCIYGRGVD350
TPEVLMYKGKGGFDKQPEIKYGDGDGTVNLAASLAALKVDSLNTVEIDGVSHTSILKDEIALKEIMKQISII420
NYELANVNAVNE

432

<210> 15
<211> 552
<212> PRT
<213> *Arabidopsis thaliana*

<400> 15
MGANSKSVTASFTVIAVFFLICGGRTAVEDETEFHGDYSKLSGIIIPGFASTQLRAWSILDCPYTPLDFN70
PLDLWLDTTKLLSAVNCWFKCMVLD PYNQTDHPECKSRPDSGLSAITELDPGYITGPLSTVWKEWLKWC140
VEFGIEANAIVAPYDWRLSPTKLEERDLYFHKLKLTFTALKLRGGPSIVFAHSMGNNVFRYFLEWLRL210
EIAPIHYLKWLDQHIHAYFAVGAPLLGSVEAIKSTLSGVTFGLPVSEGTARLLSNSFASSLWLMPPFSKNC280
KGDNTFWTHFSGGAAKDKRVYHCDEEEYQSKYSGWPTNIIINIEIPSTSARELADGTLFKAIEDYDPDSK350
RMLHQLKKYVPPFVIRNIAHRSSLAGFLLYHDDPVFNPLTPWERPPIKNVFCIYGAHLKTEVGYYFAPSG420
KPYPDNWIITDIIYETEGSLVSRSGTVVDGNAGPITGDETVPYHSLSWCKNWLGPKNITMAPQILIGKI490
KQQPEKDGSDVHVELNVDHEHGSIIANMTKAPRVKYITFYEDSESIPGKRTAVWELDKSGY 552

5

10

15

<170> PatentIn Ver. 2.0

<210> 1Q

<211> 661

20

<212> PRT

<213> Saccharomyces cerevisiae

<400> 1

25

Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30

30

Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60

35

Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95

40

Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 11045 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 140

50

Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 160Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
165 170 175

55

Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180 185 19060 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195 200 205

SUBSTITUTE SHEET (RULE 26)

Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
 210 215 220
 5 Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
 225 230 235 240
 Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
 245 250 255
 10 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
 260 265 270
 15 Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
 275 280 285
 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300
 20 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335
 25 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350
 30 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380
 35 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415
 40 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430
 45 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460
 50 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu
 485 490 495
 55 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met
 500 505 510
 60 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr
 515 520 525

Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
 530 535 540
 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
 545 550 555 560
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
 565 570 575
 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
 580 585 590
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
 595 600 605
 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
 610 615 620
 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
 625 630 635 640
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
 645 650 655
 Met Pro Phe Pro Met
 660
 <210> 2Q
 <211> 387
 <212> PRT
 <213> Arabidopsis thaliana
 <400> 2
 Val Gly Ser Asn Val Tyr Pro Leu Ile Leu Val Pro Gly Asn Gly Gly
 1 5 10 15
 Asn Gln Leu Glu Val Arg Leu Asp Arg Glu Tyr Lys Pro Ser Ser Val
 20 25 30
 Trp Cys Ser Ser Trp Leu Tyr Pro Ile His Lys Lys Ser Gly Gly Trp
 35 40 45
 Phe Arg Leu Trp Phe Asp Ala Ala Val Leu Leu Ser Pro Phe Thr Arg
 50 55 60
 Cys Phe Ser Asp Arg Met Met Leu Tyr Tyr Asp Pro Asp Leu Asp Asp
 65 70 75 80
 Tyr Gln Asn Ala Pro Gly Val Gln Thr Arg Val Pro His Phe Gly Ser
 85 90 95
 Thr Lys Ser Leu Leu Tyr Leu Asp Pro Arg Leu Arg Asp Ala Thr Ser
 100 105 110
 Tyr Met Glu His Leu Val Lys Ala Leu Glu Lys Lys Cys Gly Tyr Val
 115 120 125
 Asn Asp Gln Thr Ile Leu Gly Ala Pro Tyr Asp Phe Arg Tyr Gly Leu
 130 135 140

SUBSTITUTE SHEET (RULE 26)

Ala Ala Ser Gly His Pro Ser Arg Val Ala Ser Gln Phe Leu Gln Asp
 145 150 155 160
 5 Leu Lys Gln Leu Val Glu Lys Thr Ser Ser Glu Asn Glu Gly Lys Pro
 165 170 175
 Val Ile Leu Leu Ser His Ser Leu Gly Gly Leu Phe Val Leu His Phe
 180 185 190
 10 Leu Asn Arg Thr Thr Pro Ser Trp Arg Arg Lys Tyr Ile Lys His Phe
 195 200 205
 Val Ala Leu Ala Ala Pro Trp Gly Gly Thr Ile Ser Gln Met Lys Thr
 210 215 220
 15 Phe Ala Ser Gly Asn Thr Leu Gly Val Pro Leu Val Asn Pro Leu Leu
 225 230 235 240
 Val Arg Arg His Gln Arg Thr Ser Glu Ser Asn Gln Trp Leu Leu Pro
 245 250 255
 20 Ser Thr Lys Val Phe His Asp Arg Thr Lys Pro Leu Val Val Thr Pro
 260 265 270
 25 Gln Val Asn Tyr Thr Ala Tyr Glu Met Asp Arg Phe Phe Ala Asp Ile
 275 280 285
 Gly Phe Ser Gln Gly Val Val Pro Tyr Lys Thr Arg Val Leu Pro Leu
 290 295 300
 30 Thr Glu Glu Leu Met Thr Pro Gly Val Pro Val Thr Cys Ile Tyr Gly
 305 310 315 320
 Arg Gly Val Asp Thr Pro Glu Val Leu Met Tyr Gly Lys Gly Gly Phe
 325 330 335
 35 Asp Lys Gln Pro Glu Ile Lys Tyr Gly Asp Gly Asp Gly Thr Val Asn
 340 345 350
 40 Leu Ala Ser Leu Ala Ala Leu Lys Val Asp Ser Leu Asn Thr Val Glu
 355 360 365
 Ile Asp Gly Val Ser His Thr Ser Ile Leu Lys Asp Glu Ile Ala Leu
 370 375 380
 45 Lys Glu Ile
 385
 50 <210> 30
 <211> 389
 <212> FRT
 <213> Arabidopsis thaliana
 55 <400> 3
 Leu Lys Lys Glu Gly Leu Lys Ala Lys His Pro Val Val Phe Ile Pro
 1 5 10 15
 60 Gly Ile Val Thr Gly Gly Leu Glu Leu Trp Glu Gly Lys Gln Cys Ala
 20 25 30

Asp Gly Leu Phe Arg Lys Arg Leu Trp Gly Gly Thr Phe Leu Cys Trp
 35 40 45
 Val Glu His Met Ser Leu Asp Asn Glu Thr Gly Leu Asp Pro Ala Gly
 5 50 55 60
 Ile Arg Val Arg Ala Val Ser Gly Leu Val Ala Ala Asp Tyr Phe Ala
 65 70 75 80
 10 Pro Gly Tyr Phe Val Trp Ala Val Leu Ile Ala Asn Leu Ala His Ile
 85 90 95
 Gly Tyr Glu Glu Lys Asn Met Tyr Met Ala Ala Tyr Asp Trp Arg Leu
 100 105 110
 15 Ser Phe Gln Asn Thr Glu Arg Asp Gln Thr Leu Ser Arg Met Lys Ser
 115 120 125
 Asn Ile Glu Leu Met Val Ser Thr Asn Gly Gly Lys Lys Ala Val Ile
 20 130 135 140
 Val Pro His Ser Met Gly Val Leu Tyr Phe Leu His Phe Met Lys Trp
 145 150 155 160
 25 Val Glu Ala Pro Ala Pro Leu Gly Gly Gly Gly Gly Pro Asp Trp Cys
 165 170 175
 Ala Lys Tyr Ile Lys Ala Val Met Asn Ile Gly Gly Pro Phe Leu Gly
 180 185 190
 30 Val Pro Lys Ala Val Ala Gly Leu Phe Ser Ala Glu Ala Lys Asp Met
 195 200 205
 Arg Met Thr Arg Thr Trp Asp Ser Thr Met Ser Met Leu Pro Lys Gly
 35 210 215 220
 Gly Asp Thr Ile Trp Gly Gly Leu Asp Trp Ser Pro Glu Leu Pro Asn
 225 230 235 240
 40 Ala Pro Glu Met Glu Ile Tyr Ser Leu Tyr Gly Val Gly Ile Pro Thr
 245 250 255
 Glu Arg Ala Tyr Val Tyr Lys Leu Asn Gln Ser Pro Asp Ser Cys Ile
 260 265 270
 45 Pro Phe Gln Ile Phe Thr Ser Ala His Glu Glu Asp Glu Asp Ser Cys
 275 280 285
 Leu Lys Ala Gly Val Tyr Asn Val Asp Gly Asp Glu Thr Val Pro Val
 50 290 295 300
 Leu Ser Ala Gly Tyr Met Cys Ala Lys Ala Trp Arg Gly Lys Thr Arg
 305 310 315 320
 55 Phe Asn Pro Ser Gly Ile Lys Thr Tyr Ile Arg Glu Tyr Asn His Ser
 325 330 335
 Pro Pro Ala Asn Leu Leu Glu Gly Arg Gly Thr Gln Ser Gly Ala His
 340 345 350
 60 Val Asp Ile Met Gly Asn Phe Ala Leu Ile Glu Asp Ile Met Arg Val

SUBSTITUTE SHEET (RULE 26)

355 360 365

Ala Ala Gly Gly Asn Gly Ser Asp Ile Gly His Asp Gln Val His Ser
370 375 380

5 Gly Ile Phe Glu Trp
385

10 <210> 4Q
<211> 1986
<212> DNA
<213> *Saccharomyces cerevisiae*

15 <220>
<221> CDS
<222> (1)..(1983)

20 <400> 4
atg ggc aca ctg ttt cga aga aat gtc cag aac caa aag agt gat tct 48
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15

25 gat gaa aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga 96
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30

30 aac cac att cat cat caa cag gga tta ggc cat aag aga aga agc ggt. 144
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45

35 att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agg 192
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60

40 aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg 240
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80

45 att ttc att ctt ggt gca ttc tta ggt gta ctt ttg cct ttt agc ttt 288
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95

50 ggc gct tat cat gtt cat aat agc gat agc gac ttg ttt gac aac ttt 336
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 110

55 gta aat ttt gat tca ctt aaa gtg tat ttg gat gat tgg aaa gat gtt 384
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125

60 ctc cca caa ggt ata agt tgg ttt att gat gat att cag gct ggt aac 432
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 140

65 ttc ttc aca tct tct tta gat gat ctc agt gaa aat ttt gcc gtt ggt 480
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 160

70 aaa caa ctc tta cgt gat tat aat att gag gcc aaa cat cct gtt gta 528
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val

28/53

		165		170		175	
5	atg gtc cct ggt gtc att tct acg gga att gaa agc tgg gga gtt att						576
	Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile						
		180		185		190	
10	gga gac gat gag tgc gat agt tct gcg cat ttt cgt aaa cgg ctg tgg						624
	Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp						
		195		200		205	
15	gga agt ttt tac atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg						672
	Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp						
		210		215		220	
20	ttg aaa cat gta atg tta gat cct gaa aca ggt ctg gac cca ccg aac						720
	Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn						
		225		230		235	240
25	ttt acg cta cgt gca gca cag gcc ttc gaa tca act gat tat ttc atc						768
	Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile						
		245		250		255	
30	gca ggg tat tgg att tgg aac aaa gtt ttc caa aat ctg gga gta att						816
	Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile						
		260		265		270	
35	ggc tat gaa ccc aat aaa atg acg agt gct gcg tat gat tgg agg ctt						864
	Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu						
		275		280		285	
40	gca tat tta gat cta gaa aga cgc gat agg tac ttt acg aag cta aag						912
	Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys						
		290		295		300	
45	gaa caa atc gaa ctg ttt cat caa ttg agt ggt gaa aaa gtt tgt tta						960
	Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu						
		305		310		315	320
50	att gga cat tct atg ggt tct cag att atc ttt tac ttt atg aaa tgg						1008
	Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp						
		325		330		335	
55	gtc gag gct gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt						1056
	Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val						
		340		345		350	
60	aac gaa cac ata gat tca ttc att aat gca gca ggg acg ctt ctg ggc						1104
	Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly						
		355		360		365	
65	gct cca aag gca gtt cca gct cta att agt ggt gaa atg aaa gat acc						1152
	Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr						
		370		375		380	
70	att caa tta aat acg tta gcc atg tat ggt tgg gaa aag ttc ttc tca						1200
	Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser						
		385		390		395	400
75	aga att gag aga gta aaa atg tta caa acg tgg ggt ggt ata cca tca						1248
	Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser						
		405		410		415	

SUBSTITUTE SHEET (RULE 26)

	atg cta cca aag gga gaa gag gtc att tgg ggg gat atg aag tca tct	1296
	Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser	
	420 425 430	
5	tca gag gat gca ttg aat aac aac act gac aca tac ggc aat ttc att	1344
	Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile	
	435 440 445	
10	cga ttt gaa agg aat acg agc gat gct ttc aac aaa aat ttg aca atg	1392
	Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met	
	450 455 460	
15	aaa gac gcc att aac atg aca tta tct ata tca cct gaa tgg ctg caa	1440
	Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln	
	465 470 475 480	
20	aga aga gta cat gag cag tac tct ttc ggc tat tcc aag aat gaa gaa	1488
	Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu	
	485 490 495	
	gag tta aga aaa aat gag cta cac cac aag cac tgg tct aat cca atg	1536
	Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met	
	500 505 510	
25	gaa gta cca ctt cca gaa gct ccc cac atg aaa atc tat tgt ata tac	1584
	Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr	
	515 520 525	
30	ggc gtg aac aac cca act gaa agg gca tat gta tat aag gaa gag gat	1632
	Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp	
	530 535 540	
35	gac tcc tct gct ctg aat ttg acc atc gac tac gaa agc aag caa cct	1680
	Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro	
	545 550 555 560	
40	gta ttc ctg acc gag ggg gac gga acc gtt ccg ctg gtg gcg cat tca	1728
	Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser	
	565 570 575	
	atg tgt cac aaa tgg gcc cag ggt gct tca ccg tac aac cct gcc gga	1776
	Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly	
	580 585 590	
45	att aac gtt act att gtg gaa atg aaa cac cag cca gat cga ttt gat	1824
	Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp	
	595 600 605	
50	ata cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctg ggc agc	1872
	Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser	
	610 615 620	
55	ggc gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat	1920
	Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp	
	625 630 635 640	
60	ctc gta gag cca cgc caa ttg tct aat ttg agc cag tgg gtt tct cag	1968
	Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln	
	645 650 655	

1986

atg ccc ttc cca atg taa
Met Pro Phe Pro Met
660

5

<210> 5 a
<211> 661
<212> PRT
<213> Saccharomyces cerevisiae

10

<400> 5
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15

15

Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30

Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45

20

Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60

25

Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80

Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95

30

Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 110

Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
115 120 125

35

Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
130 135 140

40

Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
145 150 155 160

Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
165 170 175

45

Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
180 185 190

Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
195 200 205

50

Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
210 215 220

55

Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
225 230 235 240

Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
245 250 255

60

Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
260 265 270

SUBSTITUTE SHEET (RULE 26)

Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
 275 280 285
 5 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320
 10 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350
 15 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365
 20 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400
 25 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415
 30 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445
 35 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480
 40 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu
 485 490 495
 45 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met
 500 505 510
 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr
 515 520 525
 50 Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
 530 535 540
 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
 545 550 555 560
 55 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
 565 570 575
 60 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
 580 585 590

Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
595 600 605

5 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
610 615 620

Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
625 630 635 640

10 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
645 650 655

Met Pro Phe Pro Met
660

15

SEQUENCE LISTING

<110> Symme Dr., Sten
<120>

<130>
<140>
<141>
<150>

<210> 1
<211> 1986
<212> genomic DNA
<213> Saccharomyces cerevisiae
<220>
<221> CDS
<222> (1) .. (1923)

<400> 1
atg ggc aac ctg ttc cga aga aat gtc cag aac caa aag agt gat tct 48
Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
1 5 10 15
gat gcc aac aat aaa ggg ggt tct gtt cat aac aag cga gag agc aga 96
Arg Gln Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
20 25 30
aac cac att cat cat caa cag gga tta ggc cat aag aga aga agc ggt 144
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
35 40 45
att agt ggc agt gca aaa aga aat gag cgt ggc aaa gat ttc gac agc 192
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
50 55 60
aaa aga gac ggg aac ggt aga aaa cgt tgg aga gat tcc aga aga ctg 240
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
65 70 75 80
att tcc att cct ggt gaa ttc tta ggt gaa ttc atg cct ttt agc ttt 288
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
85 90 95
ggc gcc tat cat gtt cat aat agc gat agc gac ttc ttt gac aac ttt 336
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
100 105 110

gta aat ttt gat tca ctc aat gag cat ttg gat gat tgg aaa gat gtc 384
 Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
 115 120 125

ccc cca caa ggt ata agt cgg ttt acc gat gat att cag gcc ggt aac 432
 Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
 130 135 140

tac tcc aca tct tcc tta gat gat ctc agt gaa aat ttt gcc gat ggt 480
 Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
 145 150 155 160

aat caa ctc tta cgt gat cat aat acc gag gcc aat cat ccc gcc gta 528
 Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
 165 170 175

atg gcc ccc ggt gtc att tcc acg gga att gaa agc tgg gga gtt att 576
 Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
 180 185 190

gga gac gat gag tgc gat agt tct gag cat ttt cgt aaa ccc ctg tgg 624
 Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
 195 200 205

gga agt ttt ttc atg ctg aga aca atg gtt atg gat aaa gtt tgt tgg 672
 Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
 210 215 220

ttg aat cat gca atg tta gat cct gaa aca ggt ctg gac cca cgg aac 720
 Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
 225 230 235 240

ttt acg cta cgt gca gca cag ggc ttc gaa tca act gat cat ttc atc 768
 Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
 245 250 255

gca ggg cat tgg att tgg aac aat gtt ttc cca aat ctg gga gta att 816
 Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
 260 265 270

ggc cat gaa ccc aat aat atg atg agt gct ggc cat gat tgg aag ctt 864
 Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
 275 280 285

gca tac tta gat cca gaa aga cgc gat ags tac ttt acg ang cta aag 912
 Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
 290 295 300

gaa caa atc gaa ctg ttt cat caa ctg agt ggt gaa aaa gtt tgt tta 960
 Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
 305 310 315 320

att gga cat tct atg ggt tct cag att acc ttt tac ttt atg aaa tgg 1008
 Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335

gcc gag gcc gaa ggc cct ctt tac ggt aat ggt ggt cgt ggc tgg gtt 1056
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350

aac gaa cac ata gat tca ttc atc aat gca gca ggg acg ctt ctg ggc 1104
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365

gcc cca aag gca gtt cca gcc cta att agt ggt gaa atg aaa gat acc 1152
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380

att caa tta aat acg tta gcc atg tat ggt ttc gaa aag ttc ttc tca 1200
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400

aga att gag aga gca aaa atg tta caa aag tgg ggt ggt ata cca tca 1248
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415

atg cta cca aag gga gaa gag gcc att tgg ggg gat atg aag tca tct 1296
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430

tca gag gat gca ttc ant aac aac act gac aca tac ggc aat ttc att 1344
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445

cga ttt gag ags aat acg ago gat gct ttc aac aaa aat ttc aca atg 1392
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460

aat gag gcc ttt aac atg aca tta ttc aat tca cct gaa tgg ctg caa 1440
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480

aga agc gta cat gag cag tac tcc ttc ggc tat tcc aag aat gaa gaa	1488
arg arg val his glu gln tyr ser phe gly tyr ser lys asn glu glu	
485 490 495	
gag tta aga aaa aat gag cta cac cac aag cac tgg tcc aat cca atg	1536
glu leu arg lys asn glu leu his his lys his trp ser asn pro met	
500 505 510	
gaa gta cca ctt cca gaa gcc ccc cac atg aaa atc tat tgt ata tac	1584
glu val pro leu pro glu ala pro his met lys ile tyr cys ile tyr	
515 520 525	
ggg gtc aac aac cca act gaa agc gcc tat gta tat aag gaa gag gat	1632
gly val asn asn pro thr glu arg ala tyr val tyr lys glu glu asp	
530 535 540	
gac tcc tct gcc ctg aat ttg acc atc gac tac gaa agc aag caa cct	1680
asp ser ser ala leu asn leu thr ile asp tyr glu ser lys gln pro	
545 550 555 560	
gaa ttc ctg acc gag ggg gac gga acc gtt ccg ctg gtc ggc cat tca	1728
val phe leu thr glu gly asp gly thr val pro leu val ala his ser	
565 570 575	
atg tgt cac aaa cgg gcc cag ggt gcc tca ccg tac aac cct gcc gga	1776
met cys his lys trp ala gln gly ala ser pro tyr asn pro ala gly	
580 585 590	
att aac gtt act att gtc gaa atg aaa cac cag cca gat cga ttt gat	1824
ile asn val thr ile val glu met lys his gln pro asp arg phe asp	
595 600 605	
aca cgt ggt gga gca aaa agc gcc gaa cac gta gac atc ctg ggc agc	1872
ile arg gly gly ala lys ser ala glu his val asp ile leu gly ser	
610 615 620	
ggg gag ttg aac gat tac atc ttg aaa att gca agc ggt aat ggc gat	1920
ala glu leu asn asp tyr ile leu lys ile ala ser gly asn gly asp	
625 630 635 640	
ctc gcc gag cca cgc caa ttg tct aat ctg agc cag tgg gtt tct cag	1968
leu val glu pro arg gln leu ser asn leu ser gln trp val ser gln	
645 650 655	

SUBSTITUTE SHEET (RULE 26)

WO 00/60095

37/53

PCT/EP00/02701

atg ccc ttc cca atg taa
Met Pro Phe Pro Met
660

1986

SUBSTITUTE SHEET (RULE 26)

<210> 2

<211> 661

<212> PRT

<213> *Saccharomyces cerevisiae*

<400> 2

```

Met Gly Thr Leu Phe Arg Arg Asn Val Gln Asn Gln Lys Ser Asp Ser
  1          5          10          15
Asp Glu Asn Asn Lys Gly Gly Ser Val His Asn Lys Arg Glu Ser Arg
  20          25          30
Asn His Ile His His Gln Gln Gly Leu Gly His Lys Arg Arg Arg Gly
  35          40          45
Ile Ser Gly Ser Ala Lys Arg Asn Glu Arg Gly Lys Asp Phe Asp Arg
  50          55          60
Lys Arg Asp Gly Asn Gly Arg Lys Arg Trp Arg Asp Ser Arg Arg Leu
  65          70          75
Ile Phe Ile Leu Gly Ala Phe Leu Gly Val Leu Leu Pro Phe Ser Phe
  80          85          90
Gly Ala Tyr His Val His Asn Ser Asp Ser Asp Leu Phe Asp Asn Phe
  95          100          105
Val Asn Phe Asp Ser Leu Lys Val Tyr Leu Asp Asp Trp Lys Asp Val
  110          115          120
Leu Pro Gln Gly Ile Ser Ser Phe Ile Asp Asp Ile Gln Ala Gly Asn
  125          130          135
Tyr Ser Thr Ser Ser Leu Asp Asp Leu Ser Glu Asn Phe Ala Val Gly
  140          145          150
Lys Gln Leu Leu Arg Asp Tyr Asn Ile Glu Ala Lys His Pro Val Val
  155          160          165
Met Val Pro Gly Val Ile Ser Thr Gly Ile Glu Ser Trp Gly Val Ile
  170          175          180
Gly Asp Asp Glu Cys Asp Ser Ser Ala His Phe Arg Lys Arg Leu Trp
  185          190          195
Gly Ser Phe Tyr Met Leu Arg Thr Met Val Met Asp Lys Val Cys Trp
  200          205          210
Leu Lys His Val Met Leu Asp Pro Glu Thr Gly Leu Asp Pro Pro Asn
  215          220          225
Phe Thr Leu Arg Ala Ala Gln Gly Phe Glu Ser Thr Asp Tyr Phe Ile
  230          235          240
Ala Gly Tyr Trp Ile Trp Asn Lys Val Phe Gln Asn Leu Gly Val Ile
  245          250          255
Gly Tyr Glu Pro Asn Lys Met Thr Ser Ala Ala Tyr Asp Trp Arg Leu
  260          265          270
Ala Tyr Leu Asp Leu Glu Arg Arg Asp Arg Tyr Phe Thr Lys Leu Lys
  275          280          285
Glu Gln Ile Glu Leu Phe His Gln Leu Ser Gly Glu Lys Val Cys Leu
  290          295          300
  305          310          315          320

```

Ile Gly His Ser Met Gly Ser Gln Ile Ile Phe Tyr Phe Met Lys Trp
 325 330 335
 Val Glu Ala Glu Gly Pro Leu Tyr Gly Asn Gly Gly Arg Gly Trp Val
 340 345 350
 Asn Glu His Ile Asp Ser Phe Ile Asn Ala Ala Gly Thr Leu Leu Gly
 355 360 365
 Ala Pro Lys Ala Val Pro Ala Leu Ile Ser Gly Glu Met Lys Asp Thr
 370 375 380
 Ile Gln Leu Asn Thr Leu Ala Met Tyr Gly Leu Glu Lys Phe Phe Ser
 385 390 395 400
 Arg Ile Glu Arg Val Lys Met Leu Gln Thr Trp Gly Gly Ile Pro Ser
 405 410 415
 Met Leu Pro Lys Gly Glu Glu Val Ile Trp Gly Asp Met Lys Ser Ser
 420 425 430
 Ser Glu Asp Ala Leu Asn Asn Asn Thr Asp Thr Tyr Gly Asn Phe Ile
 435 440 445
 Arg Phe Glu Arg Asn Thr Ser Asp Ala Phe Asn Lys Asn Leu Thr Met
 450 455 460
 Lys Asp Ala Ile Asn Met Thr Leu Ser Ile Ser Pro Glu Trp Leu Gln
 465 470 475 480
 Arg Arg Val His Glu Gln Tyr Ser Phe Gly Tyr Ser Lys Asn Glu Glu
 485 490 495
 Glu Leu Arg Lys Asn Glu Leu His His Lys His Trp Ser Asn Pro Met
 500 505 510
 Glu Val Pro Leu Pro Glu Ala Pro His Met Lys Ile Tyr Cys Ile Tyr
 Gly Val Asn Asn Pro Thr Glu Arg Ala Tyr Val Tyr Lys Glu Glu Asp
 530 535 540
 Asp Ser Ser Ala Leu Asn Leu Thr Ile Asp Tyr Glu Ser Lys Gln Pro
 545 550 555 560
 Val Phe Leu Thr Glu Gly Asp Gly Thr Val Pro Leu Val Ala His Ser
 565 570 575
 Met Cys His Lys Trp Ala Gln Gly Ala Ser Pro Tyr Asn Pro Ala Gly
 580 585 590
 Ile Asn Val Thr Ile Val Glu Met Lys His Gln Pro Asp Arg Phe Asp
 595 600 605
 Ile Arg Gly Gly Ala Lys Ser Ala Glu His Val Asp Ile Leu Gly Ser
 610 615 620
 Ala Glu Leu Asn Asp Tyr Ile Leu Lys Ile Ala Ser Gly Asn Gly Asp
 625 630 635 640
 Leu Val Glu Pro Arg Gln Leu Ser Asn Leu Ser Gln Trp Val Ser Gln
 645 650 655
 Met Pro Phe Pro Met
 650

<210> 3^h

<211> 2312

<212> genomic DNA

<213> Schizosaccharomyces pombe

<400> 3

ATGGCGTCTT CCAAGAAGAG CAAACTCAT AAGAAAAGA AAGAAGTCAA	50
ATCTCCTATC GACTTACCA ATTCAAAGAA ACCAACTCCG GCTTTGAGTG	100
AGCAACCTTC AGCGTCCGAA ACACATCTG TTTCAATAA ATCAAGAAA	150
TCTAAATTG GAAAAAGATT GAATTTTATA TTGGGCGCTA TTTTGGGAAT	200
ATGGCGTGCT TTTTTTTTCG CTGTTGAGA CGACAATGCT GTTTTCGACC	250
CTGCTACGTT AGATAAATTT GGAATATGC TAGGCTCTTC AGACTTGTTT	300
GATCACATTA AAGGATATTT ATCTTATAAT GTGTTTAAGG ATGCACCTTT	350
TACTACGGAC AAGCCTTCGC AGTCTCCTAG CGGAATGAA GTTCAAGTTG	400
GTCTTGATAT GTACAATGAG GGATATCGAA GTGACCATCC TGTATTATG	450
GTTCCTGGTG TTATCAGCTC AGGATTAGAA AGTTGGTCTG TTAATAATTG	500
CTCGATTCCCT TACTTTAGGA AACGTCTTTG GGGTACCTGG TCTATGCTGA	550
AGGCAATGTT CCTTGACAAG CAATGCTGGC TTGAACATTT AATGCTTGAT	600
AAAAAACC GCTTGGATCC GAAGGGAATT AAGCTGCGAG CAGCTCAGGG	650
GTTCGAGCA GCTGATTTT TTATCACGGG CTATTGGATT TGGAGTAAAG	700
TAATTGAAA CCTTCTGCA ATTGGTTATG AGCCTAATA CATGTTAAGT	750
GCTTCTTACG ATTGGCGETT ATCATATGCA AATTAGAGG AACGTGATA	800
ATATTTTCA AGTTAAAA TGTTCAATTGA GTACAGCAAC ATTGTACATA	850
AGAAAAGGT AGTGTGATT TCTCACTCCA TGGGTTCAAC GGTTACGTAC	900
TAITTTTTTA AATGGGTTGA AGCTGAGGC TACGGAAATG GTGACCGAC	950
TTGGGTTAAT GATCATATTG AAGCATTTAT AAATGTGAGT CTCGATGGTT	1000
GTTCGACTAC GTTCTAAT TTTGAATAGA TATCGGGATC TTTGATTGGA	1050
CCACCCAAA CAGTGGCAGC GCTTTTATCG GGTGAATGA AAGATACAGG	1100
TATGTAAAT ACNTAARCA TGTAAATATT TAATTTTTCG TAACCGTTT	1150
AAGTCAATT GAATCASTT TCGGTCTATG GGTAAAGCAAT AATTGTTGA	1200
GATTTGTTAC TAATTACG TTTAGTTTGG AAAAATTTT TTCCCGTTCT	1250
GAAGTATATT CAAAAATACA AATGTGCTCT ACTTTTTCTA ACTTTTAATA	1300
GAGAGCCATG ATGGTTCGCA CTATGGGAGG AGTTAGTTCT ATGCTTCCTA	1350
AAGGAGCCGA TGTGTATGG GGAATGCCA GTTGGGTAAG AATATGTGC	1400

SUBSTITUTE SHEET (RULE 26)

TGTTAATTTT TTATTATAT TTAGGCTCCA GATGATCTTA ATCAAACAA	1450
TTTTTCCAAT GGTGCAATTA TTCGATATAG AGAAGACATT GATAAGGACC	1500
ACGATGAATT TGACATAGAT GATGCATTAC AATTTTTTAA AATGTTACA	1550
GATGACGATT TTAAGTCAT GCTAGCGAAA AATTATTCCC ACGGTCTTGC	1600
TTGGACTGAA AAAGAAGTGT TAAAAATAA CGAAATGCCG TCTAATGGA	1650
TAAATCCGCT AGAAGTAAGA ACATTAAAGT TACTAATTA TACTAACCCA	1700
AATAGACTAG TCTTCCTTAT GCTCCTGATA TGAAAATTTA TTGCGTTCAC	1750
CGGGTCGGAA AACCAACTGA GAGAGGTTAT TATTATACTA ATATCCTGA	1800
GGGCAACCT GTCATTGATT CCTCGGTTAA TGATGGAACA AAAGTTGAAA	1850
ATGTGAGAGA ATTTATGTTT CAAACATTCT ATTAATCTGT TTATTAGGCT	1900
ATTGTTATGG ATGATGGTGA TGGAACTTTA CCAATATTAG CCCTTGTTTT	1950
GGTGTCCAAT AAGTTTTGGC AAACAAAAG GTTTAATCCT GCTAATACAA	2000
GTATCACAAA TTATGAAATC AAGCATGAAC CTGCTGCGTT TGATCTGAGA	2050
GGAGGACCTC CCTCGGCAGA ACACGTCGAT ATACTTGGAC ATTGAGAGCT	2100
AATGTATGT TCAATTACC TTACAATTT CTATTACTAA CTCTTGAAAT	2150
AAGGAATTA TTTTAAAGT TTCATCAGGC CATGGTGACT CGGTACCAA	2200
CCGTTATATA TCAGATATCC AGTACGGACA TAAGTTTGT AGATTGCAAT	2250
TACTAACTA ACCGAACAGG GAATAATAA ATGAGATAAA TCICGATAA	2300
CCTAGAAAT AA	2312

<210> 4
<211> 3685
<212> genomic DNA
<213> Arabidopsis thaliana
<400> 4

ATGCCCTTA TTCATCGGAA AAAGCCGACG GAGAAACCAT CGACGCCGCC	50
ATCTGAAGAG GTGGTGCACG ATGAGGATTC GCAAAAGAAA CCACACGAAT	100
CTTCCAAATC CCACCATAAG AAATCGAACG GAGGAGGGAA GTGGTCGTGC	150
ATCGATTCTT GTTGTGGTGT CATTGGGTGT GTGTGTGTAA CCTGGTGGTT	200
TCTTCTCTTC CTTTACAACG CAATGCCCTGC GAGCTTCCCT CAGTATGTAA	250
CGGAGCGAAT CACGGGTCTT TTGCTGACC CGCCCGGTGT TAAGCTCAA	300
AAAGAAGGTC TTAAAGGCGA ACATCCTGTT GTCTTCATTC CTGGGATTGT	350
CACCGGTGGG CTCGAGCTTT GGAAGGCCAA ACAATGCCGT GATGGTTTAT	400
TTAGAAAACG TTTGTGGCGT GGAACTTTTG GTGAAGTCTA CAAAGGTGA	450
GCTCAACAAAT TCTCACTCTT CCTTTATATT GGGATTGGGA TTGGATCTGA	500
TGAGATCAGG CACTTGTTCG TTCTTCAACA TCACTCAACG TTTAATTCCA	550
TGTTTGTCTG TCTTACTCTT TACTTTTTTT TTTTTTTGAT GTGAAACGCT	600
ATTTTCTTAA GAGACTATTT CTGTAIGTGT AAGGTAAGCG TTCCAAGGAC	650
GTAATGGCT TGGACTATTT CTGTTTGATT GTTAACTTTA GGATATAAAA	700
TAGCTGCCCT GGAATTCGAA GTCATCTTAT TGCCAAATCT GTTGCTAGAC	750
ATGCCCTAGA GTCCGTTTAT AACAAAGTTAC TTCCTTTACT GTCGTTGCGT	800
GTAGATTAG CTTTGTGTAG CGTATAATGA AGTAGTGTTC TATGTTTTGT	850
TGGGAATAGA GAAGTTCTAA CTACATCTGT GGAAAGTGTG TTCAGGCTGT	900
GATAGAGGAC TGTTGCTTTA TTATTCAACT ATGTATATGT GTAATTAAAG	950
CTAGTTCCTT TTTGATCTTT CAGCTCAATG TGCTTTTCTC AATTTTTTTC	1000
TCAATTCGAA AGTTTCACAT CGAGTTTATT CACATGTCTT GAATTCGTC	1050
CATCCTCGTT CTTTATCCA GCTTTGAACT CCTCCCGACC CTGCTATGGA	1100
TATATTAAAA AAAAGTGTG TTGTGGGTG CATCTTTGTT ACGATCTGCA	1150
TCTTCTCTTT TCGGCTCAGT GTTCATGTTT TTGCTATGGT AGAGATGGGC	1200
AATGTTATTG TTGATGGTAA CAGTGGTATA GTTGATAGTA TCTTAACTAA	1250
TCAATTATCT CTTTGATCA GGCCTCTATG TTGGGTGGAA CAGTGTGAC	1300
TTGACAATGA AACTGGGTG GATCCAGCTG GTATTAGAGT TCGAGCTGTA	1350

TCAGGACTCG TGGCTGCTGA CTACTTTGCT CCTGGCTACT TTGTCTGGGC	1400.
AGTGCTGATT CCTAACCTTG CACATATTGG ATATGAAGAG AAAAATATGT	1450
ACATGGCTGC ATATGACTGG CGGCTTTTCTT TTCAGAACAC AGAGGTTCTT	1500
TTCTCATCGT TCTTTCTATT ATTCTGTTCC ATGTTACGTT TCTTTCTTCA	1550
TTACTTAAGG CTTAATATG TTTTCATGTTG AATTAATAGG TACGTGATCA	1600
GACTCTTAGC CGTATGAAAA GTAATATAGA GTTGATGGTT TCTACCAACG	1650
GTGGAAAAAA AGCAGTTATA GTTCCGCATT CCATGGGGGT CTGTATTTT	1700
CTACATTTTA TGAAGTGGGT TGAGGCACCA GCTCCTCTGG GTGGCGGGGG	1750
TGGGCCAGAT TGGTGTGCAA AGTATATTAA GCGGTGATG AACATTGGTG	1800
GACCATTTCT TGGTGTTCOA AAAGCTGTTG CAGGGCTTTT CTCTGCTGAA	1850
GCAAAGGATG TTGCAGTTGC CAGGTATTGA ATATCTGCTT ATACTTTTGA	1900
TGATCACAAC CTTCGGCTCTG GAACTCAAAG TTATTCTACT AAATATCAAT	1950
TCTAATAACA TTGCTATATT ATCGCTGCAA CTGACATTGG TTGATTATTT	2000
TTGCTGCTTA TGTAAGTGAA ACTCTCTTGA GATTAGACAA ATGATGAATT	2050
GATAATTCTT ACCCATTTGCT CTGTGATGAC CAGTTTCTTA GCTTCGACGA	2100
TAACATTTGT CATACTGTCT TTTGGAGGGC ATTGAATTTT GCTATGGAAA	2150
GCGCTGGAGC TTCCATGCTT GCATTCTTTA CCAATTAGCG TTATTCTGCT	2200
TCTTTCAATT TTCTTGATTA TGCATCTATG GTCTTTTATT TCTTCTTAAT	2250
TAAAGACTCG TTGGATTAGT TGCTCTATTA GTCACTTGGT TCCTTAATAT	2300
AGACTTTTAC TTTCTTCGAA AATTGCAGAG CGATTGCCCC AGGATTCTTA	2350
GACACCGATA TATTTAGACT TCAGACCTTG CAGCATGTAA TGAGATGAC	2400
ACGCACATGG GACTCAACAA TGTCTATGTT ACCGAAGGGA GGTGACACGA	2450
TATGGGGCGG GCTTGATTGG TCACCGGAGA AAGGCCACAC CTGTTGTGGG	2500
AAAAAGCATA AGAACACGA AACTTGTGGT GAAGCAGGTG AAAACGGAGT	2550
TTCAAGAAA AGTCTGTTA ACTATGGAAG GATGATATCT TTTGGGAAG	2600
AAGTAGCAGA CGCTGCGCCA TCTGAGATTA ATAATATTGA TTTTCGAGTA	2650
AGGACATATA AATCATAATA AACCTTGTAC ATTTTGTGAT TGTATGATGA	2700
ATATCTGTAC ATTTTATCTG GTGAAGGGTG CTGTCAAAGG TCAGAGTATC	2750
CGAATCACA CCTGTCTGA CGTGTGGACA GAGTACCATG ACATGGGAAT	2800
TGCTGGGATC AAGCTATCG CTGASTATAA GGTCTACACT GCTGGTGAAG	2850

CTATAGATCT ACTACATTAT GTTGCTCCTA AGATGATGGC GCGTGGTGCC	2900
GCTCATTCT CTTATGGANT TGCTGATGAT TTGGATGACA CCAAGTATCA	2950
AGATCCCAA TACTGGTCAA ATCCGTTAGA GACAAAGTAA GTGATTTCTT	3000
GATCCAACT GTATCCTTCG TCCTGATGCA TTATCAGTCT TTTTGTTC	3050
GGTCTTGTTG GATAIGGTTT TCAGCTCAA GCTTACAAAG CTGTTTCTGA	3100
GCCTTTCTCA AAAAGGCTTG CTCAGTAATA TTGAGGTGCT AAAGTTGATA	3150
CATGTGACTC TTGCTTATAA ATCCTCCGT TGSTTTGTTC TGCTTTTCA	3200
GATTACCGAA TGCTCCTGAG ATGGAAATCT ACTCTTATA CGGAGTGGG	3250
ATACCAACGG AACGAGCATA CGTATACAAG CTTAACCACT CTCCCGACAG	3300
TTGCATCCCC TTTCAGATAT TCCTTCTGC TCACGAGGAG GACGAAGATA	3350
GCTGTCTGAA AGCAGGAGTT TACAATGTGG ATGCGGATGA AACASTACCC	3400
GTCTAAGTG CCGGGTACAT GTGTGCAAA GCGTGGCGTG GCAGACAAG	3450
ATTCAACCT TCCGGAATCH AGACTTATAT AAGAGAATAC AATCACTCTC	3500
CGCCGGCTAA CCTGTTGGAA GGGCGCGGGA CGCAGAGTGG TGCCCATGTT	3550
GATATCTGG GAACTTTGC TTTGATCGAA GATATCATGA GGGTTGCCGC	3600
CGGAGGTAAC GGGTCTGATA TAGGACATGA CCAGGTCCAC TCTGGCATAT	3650
TGAAATGGTC GGAGCGTATT GACCTGAAGC TGTGA	3685

<210> 5
<211> 402
<212> cDNA
<213> Arabidopsis thaliana
<220>
<221> CDS
<222> (120)..(402)
<400> 5

```
AGAAACAGCTCTTGTCTCT CTCGACTGATCTAACAATCC CTAATCTGTGTTCTAAATTC 60
CTGGACGAGATTTGACAAAG TCCGTATAGCTTAACCTGGT TTAATTTCAAGTGACAGAT 119
ATG CCC CTT ATT CAT CGG AAA AAG CCG ACG CAG AAA CCA TCG ACG CCG 167
Met Pro Leu Ile His Arg Lys Lys Pro Thr Glu Lys Pro Ser Thr Pro
CCA TCT GAA GAG GTG GTG CAC GAT GAG GAT TCG CAA AAG AAA CCA CAC 215
Pro Ser Glu Glu Val Val His Asp Glu Asp Ser Gln Lys Lys Pro His
GAA TCT TCC AAA TCC CAC CAT AAG NAA TCG AAC GGA GGA GGG AAG TGG 263
Glu Ser Ser Lys Ser His His Lys ??? Ser Asn Gly Gly Gly Lys Trp
TCG TGC ATC GAT TCT TGT TGT TGG TTC ATT GGG TGT GTG TGT GTA ACC 311
Ser Cys Ile Asp Ser Cys Cys Trp Phe Ile Gly Cys Val Cys Val Thr
TGG TGG TTT CTT CTC TTC CTT TAC AAC GCA ATG CCT GCG AGC TTC CCT 359
Trp Trp Phe Leu Leu Phe Leu Tyr Asn Ala Met Pro Ala Ser Phe Pro
CAG TAT GTA ACG GAG CCG AAT CAC GNG TCC TTT GCC TTA CCC G 402
Gln Tyr Val Thr Glu Pro Asn His ??? Ser Phe Ala Leu Pro
```

46/53

<210> 6b
 <211> 643
 <212> cDNA
 <213> Zea mays
 <220>
 <221> CDS
 <222> (1) .. (402)
 <400> 6

```

CGG GAG AAA ATA GCT GCT TTG AAG GGG GGT GTT TAC TTA GCC GAT GGT 48
Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
1 5 10 15

GAT GAA ACT GTT CCA GTT CTT AGT GCG GGC TAC ATG TGT GCG AAA GGA 96
Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
20 25 30

TGG CGT GGC AAA ACT CCG TTC AGC CCT GCC GGC AGC AAG ACT TAC GTG 144
Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
35 40 45

AGA GAA TAC AGC CAT TCG CCA CCC TCT ACT CTC CTG GAA GGC AGG GGC 192
Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly
50 55 60

ACC CAG AGC GGT GCA CAT GTT GAT ATA ATG GGG AAC TTT GCT CTA ATT 240
Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
65 70 75 80

GAG GAC GTC ATC AGA ATA GCT GCT GGC GCA ACC GGT GAG GAA ATT GGT 288
Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
85 90 95

GGC GAT CAG GTT TAT TCA GAT ATA TTC AAG TGG TCA GAG AAA ATC AAA 336
Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
100 105 110

TTC AAA TTG TAA CCTATGGGAA GTTAAAGAAG TGCCGACCCG TTTATTGCGTTCC 391
Leu Lys Leu
115

AAAGTGTCTT GCCTGAGTGC AACTCTGGAT TTTCCTTAA TATTGTAAT TTTACAGC 449

TTCATTGCTC CCTTTGTCAA ATTACATTT GACAGGACGC CAATGCGATA CGATGTTG 507

TACCGCTATT TTCAGCATTC TATATTAAAC TGTACAGGTG TAAGTTGCAT TTGCCAGC 565

TGAAATTGTC TAGTCGTTTT CTTTACGATT TAATANCAAG TGCCGGAGCA GTGCCCCA 623

AGCNAAAAAA AAAAAAAA 643

```

SUBSTITUTE SHEET (RULE 26)

47/53

(2)

<210> 7/b
 <211> 115
 <212> PRT
 <213> Zee mays
 <400> 7

Arg Glu Lys Ile Ala Ala Leu Lys Gly Gly Val Tyr Leu Ala Asp Gly
 1 5 10 15

Asp Glu Thr Val Pro Val Leu Ser Ala Gly Tyr Met Cys Ala Lys Gly
 20 25 30

Trp Arg Gly Lys Thr Arg Phe Ser Pro Ala Gly Ser Lys Thr Tyr Val
 35 40 45

Arg Glu Tyr Ser His Ser Pro Pro Ser Thr Leu Leu Glu Gly Arg Gly
 50 55 60

Thr Gln Ser Gly Ala His Val Asp Ile Met Gly Asn Phe Ala Leu Ile
 65 70 75 80

Glu Asp Val Ile Arg Ile Ala Ala Gly Ala Thr Gly Glu Glu Ile Gly
 85 90 95

Gly Asp Gln Val Tyr Ser Asp Ile Phe Lys Trp Ser Glu Lys Ile Lys
 100 105 110

Leu Lys Leu
 115

<210> 8
<211> 616
<212> cDNA
<213> Neurospora crassa
<400> 5

GGTGGCGAAG ACCGANGGCGG AAGTTGGAGG CTAACGAGAA TGACNCTCGG	50
AGATGGATCT ACCCTCTAGA GACACGACTA CCNTTGCACC CAGCCTCAAG	100
GTNTACNGTT TTTATGGGTA GGAAGCCGAC GGAGCGAGCC TACATCTATC	150
TGGCGCCCCG TCCCGGGACG ACAACGCATC TTTAGATGAC GATCGATACG	200
ACTTTGACTN AGGGGCGACAT TGACCACGGT GTGATTTTGG GCGAAGGCGA	250
TGGCAGAGTG AACCTTAAGA GTTTGGGGTA CCTGTGCAAT AAGGGGTGGA	300
AAATGAAGAG ATACAATCCT GCGGGCTCAA AAATAACCGT GGTCCAGATG	350
CCGCATGAAC CAGAACGGTT CAATCCGAGA GGAGGGCCGA ATACGGCCGA	500
CTTAATATG TAGAAAAGGT TGAAATTTAT GAAGAGTAAT TAAATACGGC	550
ACATACGTTA CTCATAGTA TGAATAATTA AAAAAAATT TTTTTCTAA	600
AAAAAAAT AAAAAA	616

<210> 9
<211> 1562
<212> genomic DNA
<213> Arabidopsis thaliana
<400> 9

ATGAAAAAA	TATCTTCACA	TTATTCGGTA	GTCATAGCGA	TACTCGTTGT	50
GGTGACGATG	ACCTCGATGT	GTCAAGCTGT	GGGTAGCAAC	GTGTACCCTT	100
TGATTCTGGT	TCCAGGAAAC	GGAGGTAACC	AGCTAGAGGT	ACGGCTGGAC	150
AGAGAATACA	AGCCAAGTAG	TGTCTGGTGT	AGCAGCTGGT	TATATCCGAT	200
TCATAAGAAG	AGTGGTGGAT	GGTTTAGGCT	ATGGTTCGAT	GCAGCAGTGT	250
TATTGTCTCC	CTTCACCAGG	TGCTTCAGCG	ATCGAATGAT	GTTGTACTAT	300
GACCGTGATT	TGGATGATTA	CCAAATGCT	CCTGGTGTCC	AACCCCGGGT	350
TCCTCATTTT	GGTTCGACCA	AATCACTTCT	ATACCTCGAC	CCTCGTCTCC	400
GGTTAGTACT	TTCCAAGATA	TATCATTTTG	GGACATTTCG	ATAATGAACA	450
AAATAGACAT	AAATTTGGGG	GATTATTGTT	ATATCAATAT	CCATTTATAT	500
GCTAGTCGGT	AATGTGAGTG	TTATGTTAGT	ATAGTTAATG	TGAGTGTTAT	550
GTGATTTTCC	ATTTTAAATG	AAGCTAGAAA	GTTGTCGTTT	AATAATGTTG	600
CTAIGTCATG	AGAATTATAA	GGACACTATG	TAAATGTAGC	TTAATAATAA	650
GGTTTGATTT	GCAGAGATGC	CACATCTTAC	ATGGAACATT	TGGTGAAAGC	700
TCTAGAGAAA	AAATGCGGGT	ATGTTAACGA	CCAAACCATC	CTAGGAGCTC	750
CATATGATTT	CAGGTACGGC	CTGGCTGCTT	CGGGCCACCC	GTCCCGTGTA	800
GCCTCACAGT	TCCTACAAGA	CCTCAAACAA	TTGGTGGAAG	AACTAGCAG	850
CGAGAACGAA	GGAAAGCCAG	TGATACTCCT	CTCCCATAGC	CTAGGAGGAC	900
TTTTCGTCCT	CCATTTCCCT	AACCGTACCA	CCCCCTTCATG	GCGCCGCAAG	950
TACATCAAA	ACTTTGTTGC	ACTCGCTGCG	CCATGGGGTG	GGACGATCTC	1000
TCAGATGAAG	ACATTTGCTT	CTGGCAACAC	ACTCGGTGTC	CCTTTAGTTA	1050
ACCTTTTGCT	GGTCAGACGG	CATCAGAGGA	CCTCCGAGAG	TAACCAATGG	1100
CTACTTCCAT	CTACCAAGT	GTTTCACGAC	AGAACTAAAC	CGCTTGTCGT	1150
AACTCCCCAG	GTTAACTACA	CAGCTTACGA	GATGGATCGG	TTTTTTGCAG	1200
ACATTTGATT	CTCACAAGGA	GTTGTGCCTT	ACAAGACAA	AGTGTTCGCT	1250

TTAACAGAGG AGCTGATGAC TCCGGGAGTG CCAGTCACTT GCATATATGG	1300
GAGAGGAGTT GATACACCGG AGGTTTTGAT GTATGGAAAA GGAGGATTCTG	1350
ATAAGCAACC AGAGATTAAAG TATGGAGATG GAGATGGGAC GGTTAATTTG	1400
GCGAGCTTAG CAGCTTTGAA AGTCGATAGC TTGAACACCG TAGAGATTGA	1450
TGGAGTTTCG CATACATCTA TACTTAAAGA CGAGATCGCA CTTAAAGAGA	1500
TTATGAAGCA GATTTCATT ATTAATTATG AATTAGCCAA TGTTAATGCC	1550
GTCAATGAAT GA	1562

<210> 10
<211> 3996
<212> genomic DNA
<213> Arabidopsis thaliana
<400> 10

ATGGGAGCGA	ATTGGAATC	AGTAACGGCT	TCCTTCACCG	TCATCGCCGT	50
TTTTTCTTG	ATTTGCGGTG	GCCGAACCTG	GGTGGAGGAT	GAGACCGAGT	100
TTACAGGCGA	CTACTCGAAG	CTATCGGCTA	TAATCATTCG	CCCAATTCGS	150
TCCAGCGAGC	TACGAGCGTG	GTCGATCCTT	GACTGTCCAT	ACACTCCGTT	200
CGACTTCAAT	CCGCTCGACC	TCGTATGGCT	AGACACCACT	AAGGTCCGTG	250
ATCTTCAATT	CCTTCGCTCC	TTATTCTGTC	GCTCGAGTCA	CTTGTTGATG	300
AATTCGAAGC	GAATATAGC	AATGAAGCAT	GTCTCGTCTC	TCTTATTGAT	350
TCGTTTATTA	GTCAACAGTG	ACGCTTCTGA	ATCTGAGTTT	AGAGTCATAT	400
AAAACAGCTG	ACTCGGCGAG	TCTTTCCCAT	CGCTTTTGGT	TCGCTAAATG	450
TAGCGCAATG	AATGTGTAAT	TAGTCTGCGC	TTTTTATTCA	ACTAGATCTG	500
CAAGTTTTTC	AGAGTGCTCA	ATAGTAGTTA	GAATATGTTA	GGTCATTTTA	550
CTTGTCGATT	GTGATTCTTT	TGGTGTGTTG	TTACTGATCG	ACGTGATGGA	600
TGGTTTACAG	CTTCTTTCTG	CTGTCAACTG	CTGGTTTAA	TGTATGGTGC	650
TAGATCCTTA	TAATCAAAAC	GACCATCCCG	AGTGTAAAGT	ACGGCCTGAC	700
AGTGGTCTTT	CAGCCATCAC	AGNATTGGAT	CCAGSTTACA	TAACAGGTAG	750
TTTCGGATTT	TTCTTTCTTT	TGAGTTTTCT	TCAATTTGAT	ATCATCTTGT	800
TGTGATATAA	TATGGCTAAG	TTCATTAAAT	TGGTCAATTT	TCAGGTCTCT	850
TTTCTACTGT	TCGGAAGAG	TGGCTTAAGT	GGTGTCTTGA	GTTTGGTATA	900
GAAGCAATG	CAATTGTGCG	TGTTCCATAC	GATTGGAGAT	TGTCACCAAC	950
CAATTGGGAA	AGCGGTGACC	TTTACTTTCA	CAAGCTCAAG	TTAGTCTCTA	1000
TCAGGCTAAT	GTCTTTTATC	TTCTCTTTT	ATGTAAGATA	AGCTAAGAGC	1050
TCTGCTCGTC	TTCTTTTTTG	CAGGTTGACC	TTTGAAGCTG	CTTTAAAACT	1100
CCGTGCGCGC	CCTTCTATAG	TATTTGCCCA	TTCAATGGGT	AATAATGTCT	1150
TCAGATACTT	TCTGGAATGG	CTGAGGCTAG	AAATTGCACC	AAACATTAT	1200
TTGAAGTGGC	TTGATCAGCA	TATCCATGCT	TATTTGCGTG	TTGGTACCGG	1250
CCTACTATCC	TTAAGTTACC	ATTTTATTTT	TTCTCTAATT	GGGGGAGTTA	1300
TGTGTGACT	TACTGGATTG	AGCTCGATAC	CTGATTTGTT	GTTGATTTAG	1350
GAGCTCCTCT	TCTTGGTTCT	GTTGAGGCA	TCAAATCTAC	TCTCTCTGGT	1400
GTAACGTTTT	GCTTCTCTGT	TTCTGAGGTG	ACCTCTGACT	TCTCTTTAGT	1450
TTTAAGTAGT	TGATATCAAC	CAGCTCTTAT	AACCTACTGG	ATTTTCTTTT	1500
TGAAGTATT	ACTTTTGTTA	ATTGAAGTGC	TGTACGCGAT	ATGGTATCTG	1550
TAGATCTTGA	AGTGCTAGTT	ATCAAAGAAC	ATATTGTGGG	TAGTATACCT	1600
GTCAGCGGCC	TTAGCTAATA	CAACCAAAAC	ACATGTACAC	TGATTTAGTT	1650
TTAGATTAT	TATGGTAGAC	TTTAAGTTGA	CAAGAAACTT	TGACTGAAAT	1700
CTTTTATTTT	TAATAGGCTA	TGATTTGTTT	ATTGAATCA	TGTGACATAT	1750
TGACATGCGC	TTCTCATGTT	TTTTGTGGG	AAGGCTTCAG	GGAAGTGCTC	1800
GGTGTGTTG	CAATTCTTTT	GCGTCGTCT	TGTGGCTTAT	GCCATTTTCA	1850
AAGAATTGCA	AGGGTGATAA	CACATTCTGG	ACGCATTTT	CTGGGGGTGC	1900
TGCNAAGAAA	GATAAGCGCG	TATACCCTG	TGATGAAGAG	GAATATCAAT	1950
CAAAATATTC	TGGCTGGCCG	ACAAATATTA	TTAACATTGA	AATTCCTTCC	2000
ACTAGCGGTT	AGACTCTGTA	TATGCAACTG	TAACACTAAC	AAAAGTTTCA	2050
CCAAGAAATG	TCACTCTCAT	ATTTCTGTT	TTTGATGTGT	ATCCATCAGT	2100
TACAGAAACA	GCTCTAGTCA	ACATGACCAG	CATGGAATGT	GGCCTTCCCA	2150
CCCTTTTGTC	TTTCTCAGCC	CGTGAAGTAG	CAGATGGGAC	TCTTTTCAAA	2200
GCAATAGAG	ACTATGACCC	AGATAGCAAG	AGGATGTTAC	ACCAGTTAA	2250
GAATACGTA	CCTTTCTTTG	TGATAAGAAA	TATTGCTCAT	CGATCATCAC	2300
TTGCTGGGCT	CTTGATCGTC	AAATTGTTTT	GTTTAAATCT	CTATATCAAT	2350
TGTTCATATG	CTTTGTCTTT	CTTACTATAA	GAACAAGTA	TAATCAGAAA	2400
CCTTATTATT	GATTATCAGT	TCTCTCTTTA	TATTATGGAA	TGTCTTTTTC	2450
GTTTACAGTT	ATGAATGCCA	AAGGGGGTAT	TTTAGTTGAT	TGATCTCTCT	2500
ATTCTCTAGT	TTGTTTTGAC	TAATAGCGTC	AAATTTGTTT	TTCTAGCAAA	2550
TCTTTGTGAA	TTATATATAA	CATGCTAAT	ATACTTTTCA	GGTTGTATCA	2600
TGATGACCT	GTTTAAATC	CTCTGACTCC	TTGGGAGAGA	CCACCTATAA	2650
AAATGTATT	TTGCATATAT	GGTGCTCATC	TAAAGACAGA	GGTATGATGC	2700
ATTCTCAATA	TCACATTATG	CGTTGACTTT	GTTATTATAT	TCCCCATTTG	2750

GTTTGCAATA	TCTTTTIGAA	TTATGATTTA	TCTTCTCCCT	TGCATCTTAT	2800
CCTATTAAAGC	GTTAAAGGTA	CTAAATGTAT	GAAGCTGTCT	GTCAATAGGT	2850
GGTTATTACT	TTGCCCAAG	TGGCAACCT	TATCCTGATA	ATTGGATCAT	2900
CACGGATATC	ATTTATGAAA	CTGAAAGTTC	CCTCGTGTCA	AGGTAATTTT	2950
CCGCAATGSC	ACAAGTAAAA	CAGGAAGGCA	AAGTCTTCTG	TATCAGTCTA	3000
GTGGCATGTT	ATCTCAGTTG	CATAAGCAAA	TTATTAACA	ACTAAAATTT	3050
AAGTACTTTT	TTATCATTCC	TTTTGAGCTT	AGTGDATGAT	CAGTGGCTTA	3100
AAGTGGGAAG	AGGTGTTGCA	TGAAACATGA	CACTTGTATC	AAAGATAACT	3150
ACCAAAACAA	AACTAACCCA	TTTCTGAATT	TCATATTATT	AGGAGTAGTC	3200
GTGCTTTTAA	AAAATTITGT	TTAAGAAACC	GAAAAACTAG	TTCATATCTT	3250
GATTGTGCAA	TATCTGCAGG	TCTGGAAGCTG	TGGTTGATGG	GAACGCTGGA	3300
CCTATAACTG	GCGATGAGAC	GGAAGCTCA	GAAAGTTGGT	TTGAAATTAT	3350
CTTCTTGCAA	ACTACTGAAG	ACTAAGATAA	TACTTGCTTC	TGGAACACTG	3400
CTTGCTATGT	TCTCTAGTAC	ACTGCAATAT	TGACTCTCCG	CTACTTTTAT	3450
TGATTATGAA	ATTGATCTCT	TATAGGTACC	CTATCATTCA	CTCTCTTGGT	3500
GCAAGNATTC	GCTCGGACCT	AAAGTTAACA	TAACAATGGC	TCCCCAGGTA	3550
CTCTTTTFTA	GTTCCCTCACC	TTATATAGAT	CAAACTTTAA	GTGTACTTTT	3600
CTGGTTATGT	GTTGATTTAC	CTCCAATTG	TTCTTTCTAA	AAATCATATA	3650
TCCTGTACT	CCTCAAGAAC	TTGTATTAAT	CTAAACGAGA	TTCTCATTGG	3700
GAAATAAATA	CAACAGCCAG	AACACGATGG	AAGCGACGTA	CATGTGGGAC	3750
TAAATGTTGA	TCATGAGCAT	GGGTCAGACA	TCATAGCTAA	CATGACAAAA	3800
GCAACAAGGG	TTAAGTACAT	AACCTTTTAT	GAAGACTCTG	AGAGCATTCC	3850
GGGGAAGAGA	ACCGCAGTCT	GCGAGCTTGA	TAAAAGTGGG	TATTAA	3896

<210> 11b
<211> 709
<212> cDNA
<213> tomato
<400> 11

CTGGGGCCAA AAGTGAACAT AACAAAGGACA CCACAGTCAG AGCATGATGT	50
TCAGATGTAC AAGTGCATCT AATATAGAG CATCAACATG GTGAAGATAT	100
CATCCCAAT ATGACAAAGT TACCTACAAT GAAGTACATA ACCTATTATG	150
AGGATTCTCA AAGTTTTCCA GGGACAAGAA CAGCAGTTTG GGAGCTTGAT	200
AAAGCAATC ACAGGAACAT TGTCAGATCT CCAGCTTTGA TGCGGGAGCT	250
GTGGCTTGAG ATGTGGCATG ATATTCATCC TGATAAAAG TCCAAGTTTG	300
TTACAAAAGG TGGTGTCTGA TCCTCACTAT TTTCTTCTAT AATGTTTGA	350
GTTTGTATTG ACATTGTAAG TATTGCAACA AAAAGCAAAG CGTGGGCCTC	400
TCAGGGATCA GGACTGCTAT TGGGATTACG GGAAAGCTCG ATGTGCATGG	450
GCTGAACATT GTGAATACAG GTTACAATAT TCAATTATTA TTTTGCAAAA	500
TATTCTCTTT TGTGTATTT AGGCCACCTT TCCCCGGTCA CAACGATGCA	550
GATATGTATT CGGGGATGTT CACCTGGGAC AGAGTTGCAG ATTGAAGAGT	600
TCTACATCTC ACATCCTGTC ACACTATGTG TGATATTTAA GAACTTTGT	650
TTGGCGGAAC AACAAAGTTTG CACAAACATT TGAAGAAGAA AGCGAAATGA	700
TTGAGAGAG	709

THIS PAGE BLANK (USPTO)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 October 2000 (12.10.2000)

PCT

(10) International Publication Number
WO 00/60095 A3

(51) International Patent Classification⁷: C12N 15/54,
9/10, 15/81, 15/82, 1/16, 5/10, A01K 67/027, C12P 7/64

(21) International Application Number: PCT/EP00/02701

(22) International Filing Date: 28 March 2000 (28.03.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99106656.4 1 April 1999 (01.04.1999) EP
99111321.8 10 June 1999 (10.06.1999) EP
60/180,687 7 February 2000 (07.02.2000) US

(71) Applicant (for all designated States except US): BASF
PLANT SCIENCE GMBH [DE/DE]; D-67056 Lud-
wigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DAHLQVIST,
Anders [SE/SE]; Hemmansvägen 2, S-244 66 Furulund
(SE). STAHL, Ulf [SE/SE]; Liljegatan 7b, S-753 24
Uppsala (SE). LENMAN, Marit [SE/SE]; Revingegatan
13a, S-223 59 Lund (SE). BANAS, Antoni [SE/PL];

Wiolinowa 14, PL-08110 Siedlce (PL). RONNE, Hans
[SE/SE]; Dirigentvägen 169, S-756 54 Uppsala (SE).
STYMNE, Sten [SE/SE]; Torrlösa 1380, S-269 90 Svalöv
(SE).

(74) Agent: FITZNER, Uwe; Lintorfer Str. 10, D-40878
Ratingen (DE).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK,
DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,
IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,
LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT,
RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA,
UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

[Continued on next page]

(54) Title: ENZYMES OF THE BIOSYNTHETIC PATHWAY FOR THE PRODUCTION OF TRIACYLGLYCEROL AND RE-
COMBINANT DNA MOLECULES ENCODING THESE ENZYMES



(57) Abstract: The present invention relates to the isolation, identification and characterization of nucleotide sequences encoding an enzyme catalysing the transfer of fatty acids from phospholipids to diacylglycerol in the biosynthetic pathway for the production of triacylglycerol, to the said enzymes and a process for the production of triacylglycerols.

WO 00/60095 A3

WO 00/60095 A3



(88) Date of publication of the international search report:
1 February 2001

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 00/02701

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N9/10 C12N15/81 C12N15/82 C12N1/16
C12N5/10 A01K67/027 C12P7/64

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01K C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EPO-Internal, WPI Data, MEDLINE, CHEM ABS Data, BIOSIS, EMBL

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	PETER VERHASSELT ET AL.: "Twelve open reading frames revealed in the 23.6kb segment flanking the centromere on the Saccharomyces cerevisiae chromosome XIV right arm" YEAST, vol. 10, no. 7, July 1994 (1994-07), pages 1355-1361, XP002112572 abstract; table 2	1-23,27
X	-& Swissprot Database Entry Yn84_Yeast Accession number P40345; 1 February 1995 XP002112574 the whole document --- -/--	1-23,27

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

17 October 2000

Date of mailing of the international search report

30/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B

INTERNATIONAL SEARCH REPORT

Internat. Application No.

PCT/EP 00/02701

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! Database Entry SPBC776, 21 January 1999 (1999-01-21) LYNE M. ET AL.: "S. pombe chromosome II cosmid c776" Database accession no. AL035263 XP002150203 the whole document</p>	1-23,27
X	<p>DATABASE EMBL 'Online! Database Entry AI398644, 10 February 1999 (1999-02-10) XP002150204 the whole document & MARY ANNE NELSON ET AL.: "Expressed sequences from conidial, mycelial, and sexual stages of Neurospora crassa " FUNGAL GENETICS AND BIOLOGY, vol. 21, 1997, pages 348-363, XP000952173</p>	1-23,27
X	<p>KEITH STOBART ET AL.: "Triacylglycerols are synthesized and utilized by transacylation reactions in microsomal preparations of developing safflower (Carthamus tinctorius L.) seeds" PLANTA, vol. 203, no. 1, 1997, pages 58-66, XP002112573 page 58, right-hand column, last paragraph -page 59, left-hand column, paragraph 1 page 63, right-hand column, paragraph 2</p>	25
A	<p>WO 98 55631 A (CALGENE LLC) 10 December 1998 (1998-12-10) page 9, line 36 -page 10, line 7 page 12, line 28 -page 13, line 18 page 14, line 34 -page 15, line 13 page 20, line 5 -page 25, line 4</p>	1-27
P,X	<p>DATABASE SWALL 'Online! Database Entry 094680, 1 May 1999 (1999-05-01) LYNE M. ET AL.: "hypothetical 69.7 kDa protein C776.14 in chromosome II" Database accession no. 094680 XP002150205 the whole document</p>	1-23,27

maternal patent family members

Internal Allocation No.

Patent document
cited in search report

Publication
date

Patent family member(s)

Publication date

WO 9855631 A

10-12-1998

CN	1266460	T
EP	1003882	A

13-09-2000
31-05-2000

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)